

DIAGONAL GENUS 5 CURVES, ELLIPTIC CURVES OVER $\mathbb{Q}(t)$, AND RATIONAL DIOPHANTINE QUINTUPLES

MICHAEL STOLL

ABSTRACT. The problem of finding all possible extensions of a given rational diophantine quadruple to a rational diophantine quintuple is equivalent to the determination of the set of rational points on a certain curve of genus 5 that can be written as an intersection of three diagonal quadrics in \mathbb{P}^4 . We discuss how one can (try to) determine the set of rational points on such a curve. We apply our approach to the original question in several cases. In particular, we show that Fermat’s diophantine quadruple $(1, 3, 8, 120)$ can be extended to a rational diophantine quintuple in only one way, namely by $777480/8288641$.

We then discuss a method that allows us to find the Mordell-Weil group of an elliptic curve E defined over the rational function field $\mathbb{Q}(t)$ when E has full $\mathbb{Q}(t)$ -rational 2-torsion. This builds on recent results of Dujella, Gusić and Tadić. We give several concrete examples to which this method can be applied. One of these results implies that there is only one extension of the diophantine quadruple $(t - 1, t + 1, 4t, 4t(4t^2 - 1))$ over $\mathbb{Q}(t)$.

1. INTRODUCTION

A *diophantine m -tuple* is an m -tuple (a_1, \dots, a_m) of distinct nonzero integers such that $a_i a_j + 1$ is a square for all $1 \leq i < j \leq m$. A *rational diophantine m -tuple* is an m -tuple of distinct nonzero rational numbers with the same property. It is known [Duj04] that there are no diophantine sextuples and that there are only finitely many diophantine quintuples; it is conjectured that no diophantine quintuple exists. There is a recent preprint by He, Togbé and Ziegler [HTZ16] that claims to prove this. On the other hand, it has been shown that there are infinitely many rational diophantine sextuples [DKMS17], but no rational diophantine septuples are known. See Andrej Dujella’s “Diophantine m -tuples” page [Duj] for more information on background and results regarding diophantine m -tuples.

Given a (rational) diophantine quadruple (a_1, a_2, a_3, a_4) , we can ask in how many ways it can be extended to a (rational) diophantine quintuple by adding a number $a_5 \notin \{0, a_1, a_2, a_3, a_4\}$ such that

$$a_i a_5 + 1 \text{ is a square for all } 1 \leq i \leq 4.$$

Replacing a_5 by x , we obtain the system of equations

$$a_1 x + 1 = u_1^2, \quad a_2 x + 1 = u_2^2, \quad a_3 x + 1 = u_3^2, \quad a_4 x + 1 = u_4^2$$

describing a curve in \mathbb{A}^5 , whose integral (or rational) points with $x \neq 0, a_1, a_2, a_3, a_4$ correspond to the solutions of our problem (more precisely, we obtain 16 points for each solution,

Date: April 16, 2018.

2010 Mathematics Subject Classification. 11D09, 11G05, 11G30, 14G05, 14H40; 11Y50, 14G25, 14G27, 14H25, 14H52.

Key words and phrases. Diophantine quintuples, rational points, elliptic curves.

depending on how we choose the signs of the square roots u_i). This curve is irreducible and smooth; it has genus 5. Eliminating x , it is given by the three quadrics

$$a_i u_4^2 - a_4 u_i^2 - (a_i - a_4) = 0, \quad i = 1, 2, 3;$$

by homogenizing these equations we obtain a description of the smooth projective model C of the curve as an intersection of three diagonal quadrics in \mathbb{P}^4 . We call such a curve a *diagonal curve of genus 5*. Note that by Faltings' theorem [Fal83], C has only finitely many rational points, hence our original problem has only finitely many solutions for any given quadruple.

So we need to figure out a way of determining the set $C(\mathbb{Q})$ of rational points on C . This is done in Section 2. Some important features that we use are the splitting of the Jacobian J of C as a product of five elliptic curves over \mathbb{Q} , up to isogeny, and the fact that J has a large rational 2-torsion subgroup, which allows us to consider many étale double coverings of C . The Prym varieties of these coverings are isogenous (over $\bar{\mathbb{Q}}$) to a product of four elliptic curves defined (in general) over a biquadratic number field. This allows us to set up various ways of applying ‘Elliptic Curve Chabauty’ [Bru03] to our situation. This improves and extends previous work by González-Jiménez [GJ15]. We then apply this general approach to the extension problem of diophantine quadruples. More precisely, we can show that a number of quadruples from the family

$$(t - 1, t + 1, 4t, 4t(4t^2 - 1))$$

can be extended in exactly one way (the “regular” extension, which exists for every quadruple).

This prompts the question whether the regular extension is the only extension that exists generically for quadruples in the family above. To answer this question, we have to study the situation over the rational function field $\mathbb{Q}(t)$. It turns out that it is sufficient to determine the group $E(\mathbb{Q}(t))$ for a specific elliptic curve E over $\mathbb{Q}(t)$. Building on earlier work by Gusić and Tadić [GT12, GT15], which in turn uses an idea of Dujella [Duj00], we develop an approach for doing this in the case when all 2-torsion points on E are defined over $\mathbb{Q}(t)$. We finally apply our method successfully to E and use the result to show that the regular extension is the only extension to a diophantine quintuple over $\mathbb{Q}(t)$. We also show that $J(\mathbb{Q}(t))$ is generated by the differences of the 32 $\mathbb{Q}(t)$ -rational points on the associated genus 5 curve C (where J is again the Jacobian of C) and determine the structure of the group.

Acknowledgments. I would like to thank Andrej Dujella for the motivating question that led to this work and also for numerous helpful exchanges related to it. I also thank Ivica Gusić for pointing out various minor errors in earlier versions of this paper. The computations were done using the Magma Computer Algebra System [BCP97].

2. DIAGONAL GENUS 5 CURVES

Let K be a field not of characteristic 2. The canonical model of a non-hyperelliptic curve C of genus 5 over K is an intersection of three quadrics in \mathbb{P}^4 . If the coordinates on \mathbb{P}^4 can be

chosen so that the quadrics are diagonal, then we say that C is a *diagonal genus 5 curve*. We denote suitable coordinates by u_0, \dots, u_4 . Assume that we can define C in \mathbb{P}^4 by

$$\sum_{j=0}^4 a_j u_j^2 = \sum_{j=0}^4 b_j u_j^2 = \sum_{j=0}^4 c_j u_j^2 = 0$$

and let M be the 3×5 matrix whose rows are (a_0, \dots, a_4) , (b_0, \dots, b_4) , (c_0, \dots, c_4) . Then the condition for the curve defined in this way to be smooth (and hence of genus 5) is that no 3×3 minor of M vanishes. This is equivalent to saying that the net of quadrics generated by the three quadrics above does not contain quadrics of rank 2 or less. There are then exactly ten quadrics of rank 3, which are obtained by eliminating two of the variables.

Any such curve C has a subgroup A isomorphic to $(\mathbb{Z}/2\mathbb{Z})^4$ in its automorphism group, which is generated by the five involutions σ_j changing the sign of u_j and leaving the other coordinates fixed; their product is the identity.

Remark 2.1. The (coarse) moduli space of diagonal genus 5 curves has dimension 2: Fixing the coordinates, any such curve is given by a 3-dimensional space of diagonal quadratic forms in five variables. The non-vanishing of the 3×3 -minors implies that we can write the equations in the form

$$u_2^2 = au_0^2 + bu_1^2, \quad u_3^2 = a'u_0^2 + b'u_1^2, \quad u_4^2 = a''u_0^2 + b''u_1^2,$$

where the vectors (a, b) , (a', b') and (a'', b'') are linearly independent in pairs. Over an algebraically closed field, we can then scale the variables to make $a = a' = a'' = b = 1$; we are left with two parameters b', b'' that are different from 0, 1 and from each other. Permuting the variables induces an action of the symmetric group S_5 on the open subset of \mathbb{P}^2 given by the pairs (b', b'') ; the quotient is the moduli space.

We note that this is consistent with the expectation derived from the observation that the locus of Jacobians of diagonal genus 5 curves in the moduli space of principally polarized abelian 5-folds of dimension 15 is contained in the intersection of the space of Jacobians of dimension 12 and the locus of abelian varieties isogenous to a product of five elliptic curves of dimension 5.

Over a non-algebraically closed field k , we have to take into account the twists coming from the action of A , which leads to a parameterization by a subset of $k^2 \times (k^\times/k^{\times 2})^4$ (modulo the action of S_5).

Eliminating u_j leads to a double cover $\pi_j: C \rightarrow F_j$, where F_j is a curve of genus 1 given as the intersection of two diagonal quadrics in \mathbb{P}^3 . Write E_j for the Jacobian elliptic curve of F_j ; then E_j has all its 2-torsion points defined over K . (This is because F_j is an intersection of diagonal quadrics, which implies that the four singular quadrics in the pencil are all defined over the ground field.) We write J for the Jacobian variety of C . The maps $C \rightarrow F_j$ induce homomorphisms $\pi_{j,*}: J \rightarrow E_j$, which we can combine into an isogeny $\varphi: J \rightarrow \prod_{j=0}^4 E_j$ (see for example [Bre97]). Pulling back divisors under π_j induces $\pi_j^*: E_j \rightarrow J$ such that the composition $\pi_{j,*} \circ \pi_j^*$ is multiplication by 2 on E_j . The compositions $\pi_{j,*} \circ \pi_i^*$ for $i \neq j$ must be zero, since generically the elliptic curves E_i and E_j are not isogenous. Combining the π_j^* into $\hat{\varphi}: \prod_{j=0}^4 E_j \rightarrow J$, $(P_j)_j \mapsto \sum_j \pi_j^*(P_j)$, then gives an isogeny in the other direction

such that $\varphi \circ \hat{\varphi}$ is multiplication by 2 on the product of the E_j ; this implies that $\hat{\varphi} \circ \varphi$ is multiplication by 2 on J .

We can describe part of the 2-torsion subgroup $J[2]$ in the following way. Eliminating u_i and u_j results in a conic Q_{ij} ; we have the following commutative diagram:

$$\begin{array}{ccc}
 & C & \\
 \pi_i \swarrow & & \searrow \pi_j \\
 F_i & & F_j \\
 & \searrow & \swarrow \\
 & Q_{ij} &
 \end{array}$$

Pulling back any point on Q_{ij} results in an effective divisor D_{ij} of degree 4 on C such that $2D_{ij}$ is in the hyperplane class. (If the point on Q_{ij} is the image of $P \in C$, then $D_{ij} = (P) + (\sigma_i(P)) + (\sigma_j(P)) + (\sigma_i\sigma_j(P))$.) The class of D_{ij} does not depend on the chosen point, since all points on Q_{ij} are linearly equivalent. So any difference $T_{\{\{i,j\},\{i',j'\}\}} = [D_{ij} - D_{i'j'}] \in J$ is in $J[2]$. Taking $[D_{01}]$ (say) as a base-point, we obtain nine generators of the form $T_{ij} = T_{\{\{0,1\},\{i,j\}\}}$ (with $\{i,j\} \neq \{0,1\}$) for our subgroup. There are some relations among these points, though. On F_i , the sum of the four divisors obtained from the Q_{ij} with $j \neq i$ is in twice the hyperplane class, which implies that

$$\sum_{j \neq i} T_{ij} = 0 \quad \text{for each } i \in \{0, \dots, 4\}.$$

Since four of these relations imply the fifth, we get a subgroup G of $J[2]$ whose \mathbb{F}_2 -dimension is at most 5. Checking a concrete example shows that G is indeed of that dimension. One can check that $G \subset \ker(\varphi)$. Conversely, one can check that $\bigoplus_j E_j[2]$ maps surjectively onto G under $\hat{\varphi}$. This implies that the kernel of $\hat{\varphi}$ is of dimension 5 and that $G = \ker \varphi$.

Since the class of a point on Q_{ij} is defined over K , so is the class of D_{ij} , and therefore G consists of K -rational points. If $C(K)$ is non-empty or if K is a number field and C has points over all completions of K , then all the conics have K -points, and we can choose the divisors D_{ij} to be defined over K .

Now assume that $K = \mathbb{Q}$ and that $P_0 \in C(\mathbb{Q})$. Usually the differences of the images under π_i of the points in the orbit of P_0 under A will generate a group of rank 1 (changing two signs of u_j with $j \neq i$ corresponds to adding a 2-torsion point, but changing one sign and taking the difference with $\pi_i(P_0)$ will usually give a point of infinite order). So the rank of each $E_i(\mathbb{Q})$ will be positive, and the rank of $J(\mathbb{Q})$ will be at least 5; in particular, standard Chabauty techniques are not applicable, since no factor of J will have rank strictly less than its dimension. The purpose of the following is to describe an approach that can possibly succeed in determining $C(\mathbb{Q})$. It is based on covering collections and ‘Elliptic Curve Chabauty’ [Bru03]. For general information on approaches for determining the set of rational points on a curve, see also the survey [Sto11]. In the case at hand, what we do follows the ideas underlying recent work by González-Jiménez [GJ15], but we introduce some simplifications and improvements.

Since by assumption P_0 is a rational point on C , we have rational points $\pi_j(P_0)$ on the curves F_j . We then obtain isomorphisms $F_j \xrightarrow{\sim} E_j$ sending $\pi_j(P_0)$ to the origin. In the following we will identify F_j and E_j via this isomorphism. Let $\iota: C \rightarrow J$ denote the embedding sending P to the class of $(P) - (P_0)$; since P_0 is rational, ι is defined over \mathbb{Q} . We can then consider the composition δ of maps

$$C(\mathbb{Q}) \xrightarrow{\iota} J(\mathbb{Q}) \xrightarrow{\varphi} \bigoplus_{j=0}^4 E_j(\mathbb{Q}) \longrightarrow \bigoplus_{j=0}^4 \text{Sel}^{(2)}(E_j/\mathbb{Q}) \longrightarrow \bigoplus_{j=0}^4 \left(\frac{\mathbb{Q}^\times}{\mathbb{Q}^{\times 2}} \oplus \frac{\mathbb{Q}^\times}{\mathbb{Q}^{\times 2}} \right) =: H.$$

Here $\text{Sel}^{(2)}(E_j/\mathbb{Q})$ denotes the 2-Selmer group of E_j over \mathbb{Q} . The last map is the direct sum of the usual maps from 2-descent on an elliptic curve with full rational 2-torsion. For any place v of \mathbb{Q} , there is a similar map, and we obtain a commutative diagram

$$\begin{array}{ccc} C(\mathbb{Q}) & \xrightarrow{\delta} & H \\ \downarrow & & \downarrow \rho_v \\ C(\mathbb{Q}_v) & \xrightarrow{\delta_v} & H_v \end{array}$$

where H_v is defined like H , but replacing \mathbb{Q} with \mathbb{Q}_v , and the vertical maps are the natural ones.

We assume that we have determined generators of each group $E_j(\mathbb{Q})$, or at least generators of a subgroup of finite odd index, so that we know the image H_0 of $\bigoplus_j E_j(\mathbb{Q})/2E_j(\mathbb{Q})$ in H . Alternatively, it may be sufficient to know the 2-Selmer groups of the E_j , which gives us an upper bound H_0 for the image of $\bigoplus_j E_j(\mathbb{Q})/2E_j(\mathbb{Q})$ in H . The map δ and its local equivalents δ_v are given by evaluating certain rational functions on C that are defined over \mathbb{Q} at a point P and then taking the square class of the result. So δ_v is locally constant in the v -adic topology, which allows us to determine its image with a finite computation for each given place v . We do this for a number of places and define H'_0 to be the subset of H_0 consisting of elements mapping into the image of δ_v for all places v that were considered. Then $\delta(C(\mathbb{Q})) \subset H'_0$, and if we are lucky, we even may have equality, since we possibly know enough points in $C(\mathbb{Q})$ to verify that δ surjects onto H'_0 . In practice, it appears that using a few primes p of good reduction (where the computation can be essentially done over \mathbb{F}_p) suffices to cut H'_0 down to the image of the known rational points on C . (What we do here is to compute (an upper bound for) the Selmer set of C with respect to $\hat{\varphi}$; compare [Sto11] or [BS09]. Note that $\text{Sel}^{(\hat{\varphi})}(J/\mathbb{Q})$ injects into the direct sum of the 2-Selmer groups of the E_j via the map induced by φ , since the elliptic curves have full rational 2-torsion and so $\bigoplus_j E_j(\mathbb{Q})[2]$ surjects onto $J(\mathbb{Q})[\varphi] = G$.)

The action of A on C induces an action on the product of the E_j that on each E_j separately is generated by the translations by 2-torsion points and a map of the form $Q \mapsto P - Q$, where $P = \pi_j(\sigma_i(P_0))$ for some fixed $i \neq j$. This translates into the action of a subgroup H_1 of H_0 (the subgroup generated by the images of the points in the A -orbit of P_0) on H_0 by translation; the subset H'_0 is a union of cosets of this subgroup. In the computation, it will be sufficient to consider one representative of each coset, since all results can be transported by the A -action to everything in the same orbit.

Each nonzero element $T \in G$ induces an isogeny $J'_T \rightarrow J$ of degree 2; we obtain J'_T by dividing the product of the E_j by the subgroup of $\ker \hat{\varphi}$ orthogonal to T under the Weil pairing between $\ker \varphi$ and $\ker \hat{\varphi}$. Pulling back the isogeny under ι , we get an étale double cover $D_T \rightarrow C$. For each $\xi \in H'_0$ we obtain a twist $D_{T,\xi} \rightarrow C$ of this double cover with the property that all points in $C(\mathbb{Q})$ whose image in H'_0 is ξ lift to rational points on $D_{T,\xi}$. Then $D_{T,\xi}$ maps into the Prym variety of the double cover $D_{T,\xi} \rightarrow C$, which is an abelian variety of dimension 4.

One can check that all but one of the nonzero elements $T \in G$ can be represented as a difference of two divisors of the form D_{ij} . If $T = [D_{ij} - D_{kl}] = [D_{ij} + D_{kl}] - L$, where L denotes the class of a hyperplane section, then we obtain a model of $D_{T,\xi}$ in the following way. Fix points $P_{ij} \in Q_{ij}$ and $P_{kl} \in Q_{kl}$, and let $\ell_{ij}(u_0, \dots, u_4)$ and $\ell_{kl}(u_0, \dots, u_4)$ be linear forms describing the tangent lines to the conics at the points (note that ℓ_{ij} will not involve u_i and u_j ; similarly for ℓ_{kl}). Then the double covering of C given by adding the equation

$$\gamma w^2 = \ell_{ij}(u_0, \dots, u_5) \ell_{kl}(u_0, \dots, u_4)$$

will be a (singular) model of $D_{T,\xi}$, where $\gamma \neq 0$ has to be chosen so that points mapping to ξ will lift. (Concretely, if $P = (v_0 : \dots : v_4) \in C(\mathbb{Q})$ maps to ξ , then we can take $\gamma = \ell_{ij}(v_0, \dots, v_4) \ell_{kl}(v_0, \dots, v_4)$ if this value is nonzero.) If there is a rational point on C and therefore on all the conics, then we can take P_{ij} and P_{kl} to be rational, and we obtain equations over \mathbb{Q} .

For such T , we can identify the Prym variety up to isogeny as the Weil restriction of an elliptic curve defined over an étale algebra of degree 4 over \mathbb{Q} (which generically is a biquadratic number field). This can be seen by choosing the points P_{ij} and P_{kl} in such a way that the linear forms ℓ_{ij} and ℓ_{kl} together involve only three of the five variables. There are two cases.

1. $T = [D_{ij} - D_{kl}]$ with i, j, k, l distinct. Let m be such that $\{i, j, k, l, m\} = \{0, \dots, 4\}$. We can write equations for the conics Q_{ij} , Q_{kl} and Q_{ik} as

$$au_k^2 = u_m^2 - bu_l^2, \quad a'u_i^2 = u_m^2 - b'u_j^2, \quad a''u_l^2 = u_m^2 - b''u_j^2.$$

The intersection of the hyperplane $u_m = \sqrt{b}u_l$ with C is twice a divisor of the form D_{ij} and the intersection of the hyperplane $u_m = \sqrt{b'}u_j$ is twice D_{kl} . (This corresponds to taking P_{ij} to be a point with $u_k = 0$ and P_{kl} with $u_i = 0$.) Let $P \in C(\mathbb{Q})$ be a point whose image in H'_0 is ξ , say $P = (v_0 : \dots : v_4)$. Then $D_{T,\xi}$ maps to the curve $F_{T,\xi}$ given by

$$a'u_l^2 = u_m^2 - b''u_j^2, \quad \gamma w^2 = (u_m - \sqrt{b}u_l)(u_m - \sqrt{b'}u_j)$$

in \mathbb{P}^3 , where $\gamma = (v_m - \sqrt{b}v_l)(v_m - \sqrt{b'}v_j)$. This is a curve of genus 1 defined over $K = \mathbb{Q}(\sqrt{b}, \sqrt{b'})$. Let $E_{T,\xi}$ be its Jacobian elliptic curve. (Since $F_{T,\xi}$ has a point over K coming from P , it is actually an elliptic curve itself.) Then Picard functoriality induces a homomorphism $R_{K/\mathbb{Q}}E_{T,\xi} \rightarrow \text{Jac}(D_{T,\xi})$ that is an isogeny onto the Prym variety.

2. $T = [D_{ij} - D_{ik}]$ with i, j, k distinct. Let l, m be such that $\{i, j, k, l, m\} = \{0, \dots, 4\}$. We can write equations for the conics Q_{jk} , Q_{ik} and Q_{ij} as

$$au_i^2 = u_l^2 - bu_m^2, \quad a'u_j^2 = u_l^2 - b'u_m^2, \quad a''u_k^2 = u_l^2 - b''u_m^2.$$

The intersection of the hyperplane $u_l = \sqrt{b'}u_m$ with C is twice a divisor of the form D_{ik} and the intersection of the hyperplane $u_l = \sqrt{b}u_m$ is twice D_{ij} . Let $P \in C(\mathbb{Q})$ be a point

whose image in H'_0 is ξ , say $P = (v_0 : \dots : v_4)$. Then $D_{T,\xi}$ maps to the curve $F_{T,\xi}$ given by

$$au_i^2 = u_l^2 - bu_m^2, \quad \gamma w^2 = (u_l - \sqrt{b'}u_m)(u_l - \sqrt{b''}u_m)$$

in \mathbb{P}^3 , where $\gamma = (v_l - \sqrt{b'}v_m)(v_l - \sqrt{b''}v_m)$. This is a curve of genus 1 defined over $K = \mathbb{Q}(\sqrt{b'}, \sqrt{b''})$. Let $E_{T,\xi}$ be its Jacobian elliptic curve. (Since $F_{T,\xi}$ has a point over K coming from P , it is actually an elliptic curve itself.) Then Picard functoriality induces a homomorphism $R_{K/\mathbb{Q}}E_{T,\xi} \rightarrow \text{Jac}(D_{T,\xi})$ that is an isogeny onto the Prym variety.

Each of the two cases covers 15 possibilities for T . Note that for each T there can be several ways of writing it as a difference of two divisors D_{ij} , and for each such representation there can be several ways of writing down a curve $F_{T,\xi}$. They will all lead to isogenous Weil restrictions, however.

More precisely, the points T occurring in Case 1 have a single representation of the form $T = [D_{ij} + D_{kl}] - L$ in the sense that $\{\{i, j\}, \{k, l\}\}$ is uniquely determined. Since the construction above depends on choosing a representative of each of $\{i, j\}$ and $\{k, l\}$, we get four different curves $F_{T,\xi}$. The points T occurring in Case 2 have two different representations as $T = [D_{ij} + D_{ik}] - L = [D_{il} + D_{im}] - L$ (this comes from one of the relations between the $[D_{ij}]$ mentioned earlier), but each representation gives rise to only one curve $F_{T,\xi}$.

From a computational point of view, this has the advantage that we can pick the representation that involves the friendliest field K (with smallest discriminant, say) or else gives us more possibilities for computing rank bounds via Selmer groups. It turns out that we actually do get different rank bounds in general, so it makes sense to look at all of the 90 possibilities obtained by making all possible choices, together with the curves 2-isogenous to them (generically, each of the elliptic curves obtained has one point of order 2 defined over its base field).

Remark 2.2. Note that in the first case, we usually obtain elliptic curves over four different biquadratic fields whose Weil restrictions are all isogenous over \mathbb{Q} to the Prym variety of the double cover. Taking the 2-isogenous curves in two out of the four cases, we obtain four elliptic curves that become all isomorphic over the compositum K' of their various fields of definition (K' is generated by four square roots over \mathbb{Q}). Let E be this curve over K' . Then E is isomorphic over K' to all its conjugates, and the Weil restrictions are all isomorphic over K' to E^4 . The Weil restrictions are all isogenous over \mathbb{Q} , but (in general) not isomorphic (as can be seen by considering their 2-torsion subgroups, whose elements are defined over different octic fields).

We now observe that we have, in each case, an elliptic curve $F_{T,\xi}$ together with morphisms $D_{T,\xi} \rightarrow F_{T,\xi} \rightarrow \mathbb{P}^1$ over K , where the morphism to \mathbb{P}^1 is given by the quotient of any two of the u coordinates involved in $F_{T,\xi}$, such that the image of any point $P \in D_{T,\xi}(\mathbb{Q})$ in $F_{T,\xi}(K)$ maps into $\mathbb{P}^1(\mathbb{Q})$.

This is the setting for the Elliptic Curve Chabauty method [Bru03], which allows us to find the set of such points in $F_{T,\xi}(K)$ when the rank of this Mordell-Weil group is strictly less than $[K : \mathbb{Q}] = 4$. We note that González-Jiménez in [GJ15] only considers the coverings corresponding to Case 2 above.

If the degree of K is less than 4, then instead of one elliptic curve over a quartic field, we have to work with two elliptic curves over a quadratic field (this is what is done in the applications in [GJ15]) or with four elliptic curves over \mathbb{Q} .

We therefore obtain the following procedure that may determine $C(\mathbb{Q})$. For simplicity, we assume that we are in the generic case where all fields K are of degree 4.

1. For each $j \in \{0, \dots, 4\}$, compute the 2-Selmer group of E_j (or even, if possible, the group $E_j(\mathbb{Q})/2E_j(\mathbb{Q})$); determine the group H_0 and its subgroup H_1 .
2. For a suitable finite set of places v of \mathbb{Q} , determine $\delta_v(C(\mathbb{Q}_v)) \subset H_v$; use this to compute $H'_0 \subset H_0$.
Verify that the set of known points in $C(\mathbb{Q})$ surjects onto H'_0 .
3. Let X be a set of representatives of H'_0/H_1 .
4. For each $\xi \in X$, do the following.
 - a. For each $T \in G$ as in cases 1 or 2 above, determine an upper bound for the rank r of $F_{T,\xi}(K)$ (where K is the quartic field as in Case 1 or 2).
If $r \geq 4$ for all such T , report failure and stop.
 - b. For some T such that $r \leq 3$, perform the Elliptic Curve Chabauty computation to find all points $P' \in F_{T,\xi}(K)$ whose image in \mathbb{P}^1 is rational.
 - c. For each P' obtained in this way, check if it lifts to a rational point P on C . Collect all points found in this way in a set S .
5. Return S ; it is a set of orbit representatives of the action of A on $C(\mathbb{Q})$.

In Step 4a, we can for example compute the 2-Selmer group of $F_{T,\xi}$. Note that the Jacobian of each $F_{T,\xi}$ has a K -rational point of order 2, so we can also compute the 2-Selmer group of the 2-isogenous curve, which in some cases gives a better bound.

To perform Step 4b, we need to find generators of a finite-index subgroup of $F_{T,\xi}(K)$. We may get some points of infinite order from the known points in $C(\mathbb{Q})$; if this does not give enough points, then it may be difficult to find the missing generators. It might be possible to use the ‘‘Selmer group Chabauty’’ method of [Sto18] in this case, however.

3. APPLICATION TO RATIONAL DIOPHANTINE QUINTUPLES

Recall that a *rational diophantine m -tuple* is an m -tuple (a_1, \dots, a_m) of distinct nonzero rational numbers such that $a_i a_j + 1$ is a square for all $1 \leq i < j \leq m$.

Assume that a rational diophantine quadruple (a_1, a_2, a_3, a_4) is given. Then a rational number z extending it to a rational diophantine quintuple must satisfy the equations

$$a_1 z + 1 = u_1^2, \quad a_2 z + 1 = u_2^2, \quad a_3 z + 1 = u_3^2, \quad a_4 z + 1 = u_4^2$$

for suitable rational numbers u_1, \dots, u_4 . To homogenize, add another variable u_0 and set $u_0 = 1$ and $a_0 = 0$. Then we have five equations $a_j z + 1 = u_j^2$. Eliminating 1 and z then results in a diagonal genus 5 curve C , and so we can hope to apply the procedure outlined in the previous section to find all possible extensions. Note that this curve C is the locus of

points $(u_0 : \dots : u_4) \in \mathbb{P}^4$ such that

$$(3.1) \quad \text{rk} \begin{pmatrix} 1 & 1 & 1 & 1 & 1 \\ a_0 & a_1 & a_2 & a_3 & a_4 \\ u_0^2 & u_1^2 & u_2^2 & u_3^2 & u_4^2 \end{pmatrix} \leq 2.$$

The ten quadrics of rank 3 containing C are then given by the 3×3 -minors of the matrix in (3.1):

$$(a_k - a_j)u_i^2 + (a_i - a_k)u_j^2 + (a_j - a_i)u_k^2 = 0 \quad \text{for } 0 \leq i < j < k \leq 4.$$

We note that C always has the point $P_0 = (1 : 1 : 1 : 1 : 1)$ (and the points in its orbit under A), which corresponds to the ‘illegal’ extension by $z = 0$. There are also two further orbits of points corresponding to

$$(3.2) \quad z = \frac{(a_1 + a_2 + a_3 + a_4)(a_1 a_2 a_3 a_4 + 1) + 2(a_1 a_2 a_3 + a_1 a_2 a_4 + a_1 a_3 a_4 + a_2 a_3 a_4) \pm 2s}{(a_1 a_2 a_3 a_4 - 1)^2},$$

where $s = \sqrt{(a_1 a_2 + 1)(a_1 a_3 + 1)(a_1 a_4 + 1)(a_2 a_3 + 1)(a_2 a_4 + 1)(a_3 a_4 + 1)}$; compare [Duj97, Theorem 1]. Generically these three orbits are distinct, but it is possible that one of the latter two gives $z = 0$. This is always the case for the one-parameter family

$$(3.3) \quad (t - 1, t + 1, 4t, 4t(2t - 1)(2t + 1))$$

of diophantine quadruples. In any case, it is an interesting question whether for any given rational diophantine quadruple these are the only possible extensions to a rational diophantine quintuple.

As in Section 2, we use $\pi_i(P_0)$ as the origin on F_i . Let (i, j, k, l, m) be a permutation of $(0, 1, 2, 3, 4)$. Then an equation for the elliptic curve E_i is given by

$$E_i: y^2 = (x + a_j a_k + a_l a_m)(x + a_j a_l + a_k a_m)(x + a_j a_m + a_k a_l)$$

and the x -coordinate of the image of $P \in C$ is given by the quotient of linear forms

$$x = \frac{N_i(u_0, \dots, u_4)}{D_i(u_0, \dots, u_4)}$$

with

$$N_i(u_0, \dots, u_4) = \begin{vmatrix} 1 & 1 & 1 & 1 \\ a_j & a_k & a_l & a_m \\ a_j^2 & a_k^2 & a_l^2 & a_m^2 \\ a_j b_j u_j & a_k b_k u_k & a_l b_l u_l & a_m b_m u_m \end{vmatrix} \quad \text{and} \quad D_i(u_0, \dots, u_4) = \begin{vmatrix} 1 & 1 & 1 & 1 \\ a_j & a_k & a_l & a_m \\ a_j^2 & a_k^2 & a_l^2 & a_m^2 \\ u_j & u_k & u_l & u_m \end{vmatrix}$$

where $b_j = a_j - a_k - a_l - a_m$ and similarly for b_k, b_l, b_m . Then

$$x + a_j a_k + a_l a_m = \frac{V}{D_i(u_0, \dots, u_4)} \left(\frac{u_j - u_k}{a_j - a_k} + \frac{u_l - u_m}{a_l - a_m} \right)$$

is one of the expressions whose square class gives a component of the map to H . Here V is the Vandermonde determinant of (a_j, a_k, a_l, a_m) . An expression for the y -coordinate is

$$y = \frac{V \cdot ((a_j - a_k)(u_j u_k - u_l u_m) + (a_k - a_l)(u_k u_l - u_j u_m) + (a_l - a_j)(u_l u_j - u_k u_m))}{D_i(u_0, \dots, u_4) \cdot ((a_j - a_k)u_l + (a_k - a_l)u_j + (a_l - a_j)u_k)}.$$

We have implemented the algorithm of Section 2 in Magma [BCP97]; see `diophtuples.magma` at [Sto17]. For Step 4b we check if the subgroup generated by points coming from the known rational points on C reaches the upper bound for the rank. We then have generators of a finite-index subgroup and can directly perform the Elliptic Curve Chabauty computation. When there is a gap between the rank of the known subgroup and the upper bound, then we would have to find additional generators. (The most common case is that the rank bound is 3, so in principle, Elliptic Curve Chabauty is possible, but the known subgroup has rank 2. Standard conjectures imply that the rank is then 3, so we are missing one generator.) This can be quite hard for the curves showing up in the computation, so we have treated this ‘gap’ case as a failure for simplicity.

We have then used our implementation on several rational diophantine quadruples taken from the one-parameter family (3.3), with $t \in \mathbb{Q}$ positive and of small height. Excluding the cases $t = 1, \frac{1}{2}, \frac{1}{3}$ and $\frac{1}{4}$, which give degenerate quadruples, we were able to show for

$$t \in \left\{ 2, 3, \frac{2}{3}, \frac{3}{2}, 4, \frac{3}{4}, \frac{4}{3}, 5, \frac{1}{5}, \frac{2}{5}, \frac{3}{5}, \frac{5}{4}, \frac{4}{5} \right\}$$

that the extension $z \neq 0$ given by (3.2) is the only possibility. The case $t = 2$ is Fermat’s quadruple; this case is dealt with in detail (and using a variant of the approach from Section 2) in Section 4 below. When $t = \frac{3}{5}$, there is another ‘illegal’ extension by $z = \frac{12}{5}$ (which is already in the original quadruple); this is because $(\frac{12}{5})^2 + 1 = (\frac{13}{5})^2$.

In the following table, we give some more detailed information. This can be checked using file `diophtuples-verify.magma` at [Sto17].

t	T	2-isog. curve	field	subject to
2	$[D_{12} + D_{13}] - L$	no	$\mathbb{Q}(\sqrt{-6}, \sqrt{-14}, \sqrt{26})$	
3	$[D_{01} + D_{02}] - L$	no	$\mathbb{Q}(\sqrt{10}, \sqrt{13}, \sqrt{418})$	
$\frac{2}{3}$	$[D_{10} + D_{13}] - L$	no	$\mathbb{Q}(\sqrt{3}, \sqrt{-5}, \sqrt{-14})$	
$\frac{3}{2}$	$[D_{12} + D_{13}] - L$	no	$\mathbb{Q}(\sqrt{-7}, \sqrt{65}, \sqrt{165})$	
4	$[D_{14} + D_{23}] - L$	yes	$\mathbb{Q}(\sqrt{-55}, \sqrt{-335}, \sqrt{-465})$	GRH
$\frac{3}{4}$	$[D_{10} + D_{34}] - L$	no	$\mathbb{Q}(\sqrt{-1}, \sqrt{2}, \sqrt{15})$	
$\frac{4}{3}$	$[D_{02} + D_{13}] - L$	yes	$\mathbb{Q}(\sqrt{33}, \sqrt{105}, \sqrt{6097})$	GRH
5	$[D_{12} + D_{30}] - L$	no	$\mathbb{Q}(\sqrt{-2}, \sqrt{-35}, \sqrt{247})$	
$\frac{1}{5}$	$[D_{04} + D_{31}] - L$	yes	$\mathbb{Q}(\sqrt{-1}, \sqrt{3}, \sqrt{14})$	
$\frac{2}{5}$	$[D_{13} + D_{20}] - L$	no	$\mathbb{Q}(\sqrt{-7}, \sqrt{33}, \sqrt{-247})$	
$\frac{3}{5}$	$[D_{13} + D_{24}] - L$	no	$\mathbb{Q}(\sqrt{7}, \sqrt{-26}, \sqrt{334})$	
$\frac{5}{4}$	$[D_{13} + D_{24}] - L$	yes	$\mathbb{Q}(\sqrt{-6}, \sqrt{-19}, \sqrt{170})$	
$\frac{4}{5}$	$[D_{01} + D_{03}] - L$	no	$\mathbb{Q}(\sqrt{-2}, \sqrt{-35}, \sqrt{-3245})$	GRH

The column “ T ” gives the representation of the point $T \in G$ that was used to produce the curve $F_{T,\xi}$ according to the procedure in the previous section. In the column “2-isog. curve” we note whether the rank bound was obtained from $F_{T,\xi}$ itself or from the curve 2-isogenous to it. The column “field” lists the octic field for which we need class group and unit information during the computation of the 2-Selmer group of $F_{T,\xi}$ or its isogenous curve. Finally, we

indicate in the last column whether the result is conditional on the Generalized Riemann Hypothesis. The entry “GRH” indicates that this assumption was used to speed up the class group computation for the octic field, which would have been infeasible in reasonable time otherwise.

The rank of the subgroup of $F_{T,\xi}(K)$ generated by images of known points on C is 2 in all cases except for $t = \frac{3}{5}$, where it is 3. The upper bound deduced from the 2-Selmer group coincides with this lower bound in all cases. Except again for $t = \frac{3}{5}$, there is only one class ξ to consider; in the exceptional case, there are two (recall that we get the extra ‘illegal’ extension by $\frac{12}{5}$, so there are additional rational points on C).

4. POSSIBLE EXTENSIONS OF FERMAT’S QUADRUPLE

Fermat discovered the diophantine quadruple $(1, 3, 8, 120)$ and Euler found that it can be extended to a rational diophantine quintuple by adding the number $777480/8288641$. It seems to be an open question whether this is the only possibility. We now give a fairly detailed proof that this is indeed the case. (Note that the statement is contained in the results obtained in the previous section; it is the case $t = 2$. The proof given here is slightly different, though.) We do this using a variant of the approach described above, which replaces Step 2 by the computation of the fake 2-Selmer set of one of the genus 2 curves arising as quotients of the genus 5 curve.

Theorem 4.1. *The only way of extending the diophantine quadruple $(1, 3, 8, 120)$ to a rational diophantine quintuple $(1, 3, 8, 120, z)$ is to take*

$$z = \frac{777480}{8288641}.$$

Proof. We want to determine all nonzero rational z satisfying the following system of equations (with suitable $u_1, u_2, u_3, u_4 \in \mathbb{Q}$):

$$z + 1 = u_1^2, \quad 3z + 1 = u_2^2, \quad 8z + 1 = u_3^2, \quad 120z + 1 = u_4^2.$$

Writing $x = u_4$, we see that

$$x^2 + 119 = 120u_1^2, \quad x^2 + 39 = 40u_2^2, \quad x^2 + 14 = 15u_3^2,$$

so

$$5(x^2 + 119)(x^2 + 39)(x^2 + 14) = (600u_1u_2u_3)^2,$$

and we obtain a rational point on the hyperelliptic curve of genus 2

$$H: y^2 = 5(x^2 + 119)(x^2 + 39)(x^2 + 14).$$

A quick search finds the points

$$(\pm 1, \pm 600) \quad \text{and} \quad \left(\pm \frac{10079}{2879}, \pm \frac{22426285104600}{2879^3} \right).$$

The first quadruple of points corresponds to the degenerate solution $z = 0$, the second one gives Euler’s solution. We must show that these are the only rational points on H .

We perform a ‘two-cover descent’ on H as in [BS09]. This results in a two-element ‘fake 2-Selmer set’, whose elements are accounted for by the ‘trivial’ points in $H(\mathbb{Q})$ with $x = \pm 1$. In particular, we see that the automorphism of H given by changing the sign of x interchanges

the two Selmer set elements, so it suffices to find all rational points mapping to one of them, say the element corresponding to $x = 1$. Any such rational point will give rise to a point over $K = \mathbb{Q}(\sqrt{-119}, \sqrt{-39})$ on the curve

$$E: Y^2 = 15(1 - \sqrt{-119})(1 - \sqrt{-39}) \cdot (X^2 + 14)(X - \sqrt{-119})(X - \sqrt{-39})$$

with X -coordinate in \mathbb{Q} (since $X = x$). This curve E is an elliptic curve over K ; it has one K -rational point of order 2. We compute its 2-Selmer group and find that it is isomorphic to $(\mathbb{Z}/2\mathbb{Z})^3$. This implies that the rank of $E(K)$ is at most 2. We find two independent points in $E(K)$ (coming from the known points on H), so the rank is indeed 2. We saturate the group generated by the 2-torsion point and these two points at 2, 3, 5, 7, which is enough for the Elliptic Curve Chabauty computation as implemented in Magma [BCP97]. This computation finally shows that the only points in $E(K)$ with $X \in \mathbb{Q}$ are the points with $X = 1$ or $X = 10079/2879$. This concludes the proof. \square

Remark 4.2. Since the curve H in the proof is bi-elliptic, with both elliptic curve quotients of rank 1, it would also be possible to use ‘‘Quadratic Chabauty’’ to determine its set of rational points. See recent work by Balakrishnan and Dogra [BD, BD17].

5. MORDELL-WEIL GROUPS OF ELLIPTIC CURVES OVER $\mathbb{Q}(t)$

We now consider the problem of determining generators for the group $E(\mathbb{Q}(t))$ of $\mathbb{Q}(t)$ -rational points on an elliptic curve E defined over $\mathbb{Q}(t)$, in the case when all points of order 2 on E are defined over $\mathbb{Q}(t)$. The usual 2-descent approach gives an embedding

$$\delta: \frac{E(\mathbb{Q}(t))}{2E(\mathbb{Q}(t))} \longrightarrow H_0 \subset \frac{\mathbb{Q}(t)^\times}{\mathbb{Q}(t)^{\times 2}} \times \frac{\mathbb{Q}(t)^\times}{\mathbb{Q}(t)^{\times 2}}$$

with a suitable finite subgroup H_0 . For all but finitely many $\tau \in \mathbb{Q}$, we can specialize E to an elliptic curve E_τ over \mathbb{Q} , and we have a natural homomorphism $\rho_\tau: E(\mathbb{Q}(t)) \rightarrow E_\tau(\mathbb{Q})$. By Silverman’s specialization theorem [Sil83], ρ_τ is injective once τ has sufficiently large height. If we can find such a τ with the property that the subgroup of $E(\mathbb{Q}(t))$ generated by some known points surjects onto $E_\tau(\mathbb{Q})$, then this proves that our known points already generate $E(\mathbb{Q}(t))$. The lower bound for the height of τ that one can extract from Silverman’s approach tends to be too large to be practical; also, ρ_τ is usually injective also for most ‘‘small’’ τ . So it is useful to have a more concrete computational criterion for testing the injectivity of ρ_τ . Such a criterion was provided by Gusić and Tadić in a recent paper [GT15] (building on the earlier paper [GT12]), in the case when E has at least one point of order 2 defined over $\mathbb{Q}(t)$. We improve on their approach somewhat by making use of several specializations to cut down the group H_0 that contains the image of $E(\mathbb{Q}(t))/2E(\mathbb{Q}(t))$, which gives the criterion a better chance of success.

We then use our method to determine generators of the Mordell-Weil group of the elliptic curve

$$E: y^2 = (x + 4t(t - 1))(x + 4t(t + 1))(x + (t - 1)(t + 1))$$

over $\mathbb{Q}(t)$. It turns out that the group has rank 1 and maps isomorphically under ρ_2 to $E_2(\mathbb{Q})$. We then use this result to show that the ‘‘regular’’ extension of a family of rational diophantine quadruples to a rational diophantine quintuple is the only generic (i.e., given by a rational function in the parameter) such extension; see Sections 6 and 7 below.

So let E be an elliptic curve over the rational function field $\mathbb{Q}(t)$. Since $\mathbb{Q}(t)$ is a finitely generated field, the Mordell-Weil group $E(\mathbb{Q}(t))$ is a finitely generated abelian group [LN59]. In the following, we will assume that all points of order 2 on E are defined over $\mathbb{Q}(t)$ and that E is given by a Weierstrass equation with coefficients in $\mathbb{Z}[t]$ of the form

$$E: y^2 = (x - e_1(t))(x - e_2(t))(x - e_3(t)).$$

As over any field, we have the exact sequence

$$0 \longrightarrow E(\mathbb{Q}(t))[2] \longrightarrow E(\mathbb{Q}(t)) \xrightarrow{\cdot 2} E(\mathbb{Q}(t)) \xrightarrow{\delta} H,$$

where H is the subgroup of $(\mathbb{Q}(t)^\times/\mathbb{Q}(t)^{\times 2})^3$ consisting of triples such that the product of the three entries is trivial. One can show in the usual way that the image of δ is contained in the subgroup generated by the prime divisors in the UFD $\mathbb{Z}[t]$ of

$$\Delta(t) = (e_1(t) - e_2(t))(e_1(t) - e_3(t))(e_2(t) - e_3(t)) \in \mathbb{Z}[t]$$

in each of the three components; see [GT15].

Let $\tau \in \mathbb{Q}$ be such that $\Delta(\tau) \neq 0$. Then we can specialize E to an elliptic curve E_τ over \mathbb{Q} , and we obtain a specialization homomorphism $\rho_\tau: E(\mathbb{Q}(t)) \rightarrow E_\tau(\mathbb{Q})$. Gusić and Tadić in [GT15] give a criterion for when ρ_τ is injective. In the following, we will give a streamlined proof of their result (which is based on the same ideas), which we will then use to devise a method that can show that a known set of points in $E(\mathbb{Q}(t))$ generates the latter group.

We begin with an easy lemma.

Lemma 5.1. *Assume that we have the following commutative diagram of abelian groups with exact rows:*

$$\begin{array}{ccccccc} A & \longrightarrow & B & \xrightarrow{\varphi} & C & \longrightarrow & D \\ \alpha \downarrow & & \beta \downarrow & & \downarrow \gamma & & \downarrow \delta \\ A' & \longrightarrow & B' & \longrightarrow & C' & \longrightarrow & D' \end{array}$$

Assume further that α is surjective and δ is injective. Then φ induces a surjective homomorphism $\ker \beta \rightarrow \ker \gamma$.

Proof. This is an easy diagram chase. □

Corollary 5.2. *Let E and H be as above. Assume that $H' \subset H$ is finitely generated and contains $\delta(E(\mathbb{Q}(t)))$. If the specialization homomorphism $h_\tau: H' \rightarrow H_\tau = (\mathbb{Q}^\times/\mathbb{Q}^{\times 2})^3$ associated to τ is injective, then $\rho_\tau: E(\mathbb{Q}(t)) \rightarrow E_\tau(\mathbb{Q})$ is also injective.*

Proof. We first remark that ρ_τ is injective on torsion. Let $K_\tau = \ker \rho_\tau$; then K_τ is a finitely generated torsion-free abelian group, so K_τ is a free abelian group. We consider the commutative diagram with exact rows

$$\begin{array}{ccccccc} E(\mathbb{Q}(t))[2] & \longrightarrow & E(\mathbb{Q}(t)) & \xrightarrow{\cdot 2} & E(\mathbb{Q}(t)) & \xrightarrow{\delta} & H' \\ \simeq \downarrow & & \rho_\tau \downarrow & & \rho_\tau \downarrow & & \downarrow h_\tau \\ E_\tau(\mathbb{Q})[2] & \longrightarrow & E_\tau(\mathbb{Q}) & \xrightarrow{\cdot 2} & E_\tau(\mathbb{Q}) & \xrightarrow{\delta_\tau} & H_\tau. \end{array}$$

Since E has full 2-torsion over $\mathbb{Q}(t)$, the leftmost map is an isomorphism. By assumption the rightmost map is injective. So Lemma 5.1 tells us that $2K_\tau = K_\tau$. Since K_τ is free, this implies $K_\tau = 0$ as desired. \square

This easily implies Theorem 1.1 of [GT15]: their condition is equivalent to the injectivity of $H' \rightarrow H_\tau$, where H' is a slightly refined version of the general upper bound for $\delta(E(\mathbb{Q}(t)))$ mentioned above.

Remark 5.3. If $E(\mathbb{Q}(t))[2] \simeq \mathbb{Z}/2\mathbb{Z}$ and E' is the 2-isogenous curve with dual isogenies $\phi: E \rightarrow E'$ and $\hat{\phi}: E' \rightarrow E$, then we can use

$$\begin{array}{ccccccc} E(\mathbb{Q}(t))[2] \times E'(\mathbb{Q}(t))[2] & \longrightarrow & E(\mathbb{Q}(t)) \times E'(\mathbb{Q}(t)) & \xrightarrow{\Phi} & E(\mathbb{Q}(t)) \times E'(\mathbb{Q}(t)) & \xrightarrow{\delta} & H' \\ \downarrow & & \rho_\tau \downarrow & & \rho_\tau \downarrow & & \downarrow h_\tau \\ E_\tau(\mathbb{Q})[2] \times E'_\tau(\mathbb{Q})[2] & \longrightarrow & E_\tau(\mathbb{Q}) \times E'_\tau(\mathbb{Q}) & \xrightarrow{\Phi} & E_\tau(\mathbb{Q}) \times E'_\tau(\mathbb{Q}) & \xrightarrow{\delta_\tau} & H_\tau. \end{array}$$

with $\Phi(P, Q) = (\hat{\phi}(Q), \phi(P))$ and $H' \subset (\mathbb{Q}(t)^\times / \mathbb{Q}(t)^{\times 2})^2$ coming from the descent maps associated to ϕ and $\hat{\phi}$. Note that $\Phi \circ \Phi$ is multiplication by 2 on $E \times E'$. If the leftmost map is surjective and h_τ is injective, then both specialization maps $E(\mathbb{Q}(t)) \rightarrow E_\tau(\mathbb{Q})$ and $E'(\mathbb{Q}(t)) \rightarrow E'_\tau(\mathbb{Q})$ are injective. This recovers Theorem 1.3 of [GT15]; note that their condition implies that there are no additional rational 2-torsion points on the specialized curves.

Remark 5.4. The exact same result clearly holds when we replace E by the Jacobian variety J of a hyperelliptic curve with the property that all 2-torsion points on J are $\mathbb{Q}(t)$ -rational. Remark 5.3 generalizes in a similar way to Jacobians of curves of genus 2 admitting a Richelot isogeny whose kernel consists of points defined over $\mathbb{Q}(t)$.

Remark 5.5. If $E(\mathbb{Q}(t))[2] = 0$, then one can still use this approach in principle (one needs to use specializations with $E_\tau(\mathbb{Q})[2] = 0$); however, the difficulty lies in obtaining a suitable bounding group H' . It is a subgroup of the group of square classes in the function field of the curve given by the 2-torsion sections in the elliptic surface associated to E , and so it will very likely be necessary to obtain information on the 2-torsion in the Picard group of this curve (which is trigonal, but can be arbitrarily complicated in any other way).

If one wants to use the criterion of Gusić and Tadić directly, one needs to find τ such that -1 and the distinct prime factors of $\Delta(t)$ (in the UFD $\mathbb{Z}[t]$ and up to sign) have independent images in $\mathbb{Q}^\times / \mathbb{Q}^{\times 2}$. If there are many prime factors, this usually means that τ cannot be taken to have small height. This in turn may lead to difficulties when trying to determine the rank of $E_\tau(\mathbb{Q})$.

We suggest the following modified approach. Let $P_1, \dots, P_r \in E(\mathbb{Q}(t))$ be known independent points. We want to show that they generate the free part of $E(\mathbb{Q}(t))$.

1. Find the prime divisors of Δ in $\mathbb{Z}[t]$ and let H_0 be the subgroup of H consisting of triples all of whose entries are represented by a divisor of Δ . (Then the image of δ is contained in H_0 .)

2. For a finite set T of $\tau \in \mathbb{Q}$ such that $\Delta(\tau) \neq 0$, compute the 2-Selmer group of E_τ as a subgroup $S_\tau \subset H_\tau$ (if feasible, compute tighter bounds for $\delta_\tau(E_\tau(\mathbb{Q}))$, for example by determining $E_\tau(\mathbb{Q})$).
3. Set $H' := \{a \in H_0 : \forall \tau \in T : h_\tau(a) \in S_\tau\}$.
4. Now consider values of τ such that $\Delta(\tau) \neq 0$ and $H' \cap \ker h_\tau = 0$. If $\rho_\tau(P_1), \dots, \rho_\tau(P_r)$ generate the free part of $E_\tau(\mathbb{Q})$, then P_1, \dots, P_r generate the free part of $E(\mathbb{Q}(t))$.

To see that Step 4 works, first note that by Corollary 5.2, we know that ρ_τ is injective. If the known subgroup of the free part of $E(\mathbb{Q}(t))$ surjects under ρ_τ onto the free part of $E_\tau(\mathbb{Q})$, then it follows that the known subgroup must already be all of the free part.

We have implemented this procedure in Magma [BCP97]; see file `ellQt.magma` at [Sto17].

6. EXAMPLES

In the following, we will use additive notation for the group H and its subgroups. Note that they are killed by 2, so they can be considered as vector spaces over \mathbb{F}_2 .

We consider the curve

$$E: y^2 = (x + 4t(t-1))(x + 4t(t+1))(x + (t-1)(t+1)).$$

It has the point

$$P = (0, 4t(t-1)(t+1))$$

of infinite order. The discriminant factors as

$$\Delta_E = -2^{10}t^2(t-1)^2(t+1)^2(3t-1)^2(3t+1)^2,$$

hence the image of $E(\mathbb{Q}(t))$ under δ is contained in H_0 , where

$$H_0 = H \cap \langle -1, 2, t, t-1, t+1, 3t-1, 3t+1 \rangle^3,$$

so that H_0 has dimension 14. The image of the known subgroup of $E(\mathbb{Q}(t))$ is H_1 , generated by

$$\begin{aligned} &(-2t(t-1)(3t-1), 2t, -(t-1)(3t-1)), \\ &(-2t, 2t(t+1)(3t+1), -(t+1)(3t+1)) \quad \text{and} \\ &(t(t-1), t(t+1), (t-1)(t+1)). \end{aligned}$$

We can check that for the specializations E_t with $t = 2$, $t = 3$ and $t = 5$, the groups $E_t(\mathbb{Q})$ are generated by the specializations of the known generators of $E(\mathbb{Q}(t))$. (The Magma function `MordellWeilGroup` does this for us.) This implies that

$$H_1 \subset \delta(E(\mathbb{Q}(t))) \subset (H_1 + \ker(h_2)) \cap (H_1 + \ker(h_3)) \cap (H_1 + \ker(h_5)).$$

We can easily check that the intersection on the right equals H_1 , so that $\delta(E(\mathbb{Q}(t))) = H_1$ (this already shows that $E(\mathbb{Q}(t))$ has rank 1 and that the known points generate a subgroup of finite odd index).

For $\tau = 2$ (and also $\tau = 3$ and $\tau = 5$), we have that H_1 meets the kernel of h_τ trivially. Since the specializations of the known generators of $E(\mathbb{Q}(t))$ are generators of $E_\tau(\mathbb{Q})$, we have shown that the known points actually generate the full group $E(\mathbb{Q}(t))$:

Proposition 6.1. *The group $E(\mathbb{Q}(t))$ is isomorphic to $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}$, with generators*

$$T_1 = (-4t(t-1), 0), \quad T_2 = (-4t(t+1), 0) \quad \text{and} \quad P = (0, 4t(t-1)(t+1)).$$

Remark 6.2. The smooth elliptic surface \mathcal{E} over \mathbb{P}^1 associated to E is a rational surface. It has six bad fibers (over $t = 0, 1, -1, \frac{1}{3}, -\frac{1}{3}$ and ∞), which are all of type I_2 . By the Shioda-Tate formula [Shi99], this gives

$$\text{rank } E(\bar{\mathbb{Q}}(t)) = 10 - 2 - 6 = 2,$$

so there must be another point defined over $K(t)$ for some finite extension K of \mathbb{Q} . After searching a bit, one finds the independent point

$$Q = (2t(t+1), 2\sqrt{3}t(t+1)(3t-1)),$$

which is defined over $\mathbb{Q}(\sqrt{3})$ (and actually comes from the quadratic twist of E by 3). The height pairing matrix of P and Q is the diagonal matrix with entries $\frac{1}{2}, \frac{1}{2}$. Since the bad fibers are of type I_2 , it follows that the canonical height of any point is in $\frac{1}{2}\mathbb{Z}$. Since there is no proper superlattice of the square lattice generated by P and Q whose points satisfy this condition, this implies that P and Q generate the free part of $E(\bar{\mathbb{Q}}(t))$. We thus obtain an alternative proof of Proposition 6.1. We would also like to mention that Dujella [Duj00, Theorem 4] had already shown that $E(\mathbb{Q}(t))$ has rank 1.

In the same way, one can show the following.

Proposition 6.3.

(1) *The group of $\mathbb{Q}(t)$ -points on the elliptic curve*

$$y^2 = (x + (t^2 - 1))(x + 4t(t-1)(4t^2 - 1))(x + 4t(t+1)(4t^2 - 1))$$

is isomorphic to $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}^2$, generated by the points

$$\begin{aligned} &(- (t^2 - 1), 0), \quad (-4t(t-1)(4t^2 - 1), 0), \quad (0, 4t(t^2 - 1)(4t^2 - 1)) \\ &\text{and} \quad (-8t^2(2t^2 - 1), 4t(4t^2 - t - 1)(4t^2 + t - 1)). \end{aligned}$$

(2) *The group of $\mathbb{Q}(t)$ -points on the elliptic curve*

$$y^2 = (x + 4t(t-1))(x + 4t(t-1)(4t^2 - 1))(x + 16t^2(4t^2 - 1))$$

is isomorphic to $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}^2$, generated by the points

$$\begin{aligned} &(-4t(t-1), 0), \quad (-4t(t-1)(4t^2 - 1), 0), \quad (0, 16t^2(t-1)(4t^2 - 1)) \\ &\text{and} \quad (4t(t-1)(2t-1)(4t+1), 16t^2(t-1)(2t-1)(3t+1)(4t-1)). \end{aligned}$$

(3) *The group of $\mathbb{Q}(t)$ -points on the elliptic curve*

$$y^2 = (x + 4t(t+1))(x + 4t(t+1)(4t^2 - 1))(x + 16t^2(4t^2 - 1))$$

is isomorphic to $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}^2$, generated by the points

$$\begin{aligned} &(-4t(t+1), 0), \quad (-4t(t+1)(4t^2 - 1), 0), \quad (0, 16t^2(t+1)(4t^2 - 1)) \\ &\text{and} \quad (4t(t+1)(2t+1)(4t-1), 16t^2(t+1)(2t+1)(3t-1)(4t+1)). \end{aligned}$$

(4) *The group of $\mathbb{Q}(t)$ -points on the elliptic curve*

$$y^2 = x(x + 16t(2t^2 - 1))(x + (3t + 1)(4t + 1)(4t^2 - t - 1))$$

is isomorphic to $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}$, generated by the points

$$(0, 0), \quad (-16t(2t^2 - 1), 0) \quad \text{and} \quad (2(3t + 1), 2(3t + 1)(4t - 1)(4t^2 + t - 1)).$$

Proof. Running our implementation on the four given curves and the given points results in a proof that the points generate the group in each case. See the commented-out last part of `ellQt.magma` for the relevant commands. \square

We note that the specializations at integers of the curve considered in part (1) have been studied by Najman [Naj09].

Further applications of the method can be found in [DGT15, DPT16].

7. GENERIC EXTENSIONS OF THE FAMILY OF DIOPHANTINE QUADRUPLES

As already mentioned, there is a one-parameter family of rational diophantine quadruples, given by

$$(a_1, a_2, a_3, a_4) = (t - 1, t + 1, 4t, 4t(4t^2 - 1)) \quad \text{with } t \in \mathbb{Q} \setminus \left\{-1, -\frac{1}{2}, -\frac{1}{3}, -\frac{1}{4}, 0, \frac{1}{4}, \frac{1}{3}, \frac{1}{2}, 1\right\}.$$

We have seen in Section 3 that it can be extended to a quintuple by setting

$$a_5 = \frac{4t(2t - 1)(2t + 1)(4t^2 - 2t - 1)(4t^2 + 2t - 1)(8t^2 - 1)}{(64t^6 - 80t^4 + 16t^2 - 1)^2}$$

We can now use Proposition 6.1 to show that this extension is in fact the only *generic* extension of the quadruple, in the sense that there is no other rational function $a_5(t) \in \mathbb{Q}(t)^\times$ such that $(a_1(t), \dots, a_5(t))$ is a diophantine quintuple in $\mathbb{Q}(t)$ (i.e., such that $a_i(t)a_j(t) + 1$ is a square in $\mathbb{Q}(t)$ for all $1 \leq i < j \leq 5$):

Theorem 7.1. *The only function $f(t) \in \mathbb{Q}(t)^\times$ such that*

$$(t - 1)f(t) + 1, \quad (t + 1)f(t) + 1, \quad 4tf(t) + 1 \quad \text{and} \quad 4t(4t^2 - 1)f(t) + 1$$

are all squares in $\mathbb{Q}(t)$ is

$$f(t) = \frac{4t(2t - 1)(2t + 1)(4t^2 - 2t - 1)(4t^2 + 2t - 1)(8t^2 - 1)}{(64t^6 - 80t^4 + 16t^2 - 1)^2}.$$

Proof. Any such $f(t)$ gives rise to a $\mathbb{Q}(t)$ -rational point on the curve C defined by

$$(t - 1)x + 1 = u_1^2, \quad (t + 1)x + 1 = u_2^2, \quad 4tx + 1 = u_3^2, \quad 4t(4t^2 - 1)x + 1 = u_4^2$$

with $x = f(t)$. We have a morphism

$$\pi: C \longrightarrow E, \quad (x, u_1, u_2, u_3, u_4) \longmapsto (4t(t - 1)(t + 1)x, 4t(t - 1)(t + 1)u_1u_2u_3),$$

where E is the elliptic curve considered in Proposition 6.1. The morphism extends to a finite morphism of degree 8 from the projective closure of C to E . Now consider the following commutative diagram.

$$\begin{array}{ccc} C(\mathbb{Q}(t)) & \xrightarrow{\pi} & E(\mathbb{Q}(t)) \\ \downarrow \rho_{C,2} & & \simeq \downarrow \rho_2 \\ C_2(\mathbb{Q}) & \xrightarrow{\pi_2} & E_2(\mathbb{Q}) \end{array}$$

Here the subscript 2 denotes the specializations at $t = 2$. Since the genus of C_2 is $5 \geq 2$, there are only finitely many points in $C_2(\mathbb{Q})$. In fact, we have shown in Theorem 4.1 that there are precisely 32 points in $C_2(\mathbb{Q})$, corresponding to the extension of $(1, 3, 8, 120)$ by (the illegal value) zero and by $777480/8288641$ (recall that each extension gives rise to an A -orbit of 16 points on C_2). This implies that the image of $C_2(\mathbb{Q})$ in $E_2(\mathbb{Q})$ consists of $\{\pm P_2, \pm 5P_2\}$ (where $P_2 = (0, 24) \in E_2(\mathbb{Q})$ is the specialization of P). Here the points giving the extension by zero map to $\pm P_2$, and the points giving the nontrivial extension map to $\pm 5P_2$.

Since the right-hand vertical map is an isomorphism by Proposition 6.1, this shows that any point in $C(\mathbb{Q}(t))$ must map to $\pm P$ or $\pm 5P$ in $E(\mathbb{Q}(t))$. This leads to $x = 0$ or $x = f(t)$, finishing the proof. \square

The curve E and the other four elliptic curves showing up in Proposition 6.3 are the five elliptic curves occurring as quotients of the genus 5 curve C over $\mathbb{Q}(t)$ associated to the quadruple $(t - 1, t + 1, 4t, 4t(4t^2 - 1))$. So Propositions 6.1 and 6.3 together imply that the group $J(\mathbb{Q}(t))$ has rank 8, where J denotes the Jacobian variety of C . We can actually say more.

Proposition 7.2. *Let C denote the genus 5 curve over $\mathbb{Q}(t)$ associated to the diophantine quadruple*

$$(t - 1, t + 1, 4t, 4t(4t^2 - 1))$$

over $\mathbb{Q}(t)$, and denote its Jacobian by J . Then C has exactly 32 $\mathbb{Q}(t)$ -points, and the group $J(\mathbb{Q}(t))$ is isomorphic to $(\mathbb{Z}/2\mathbb{Z})^5 \times \mathbb{Z}^8$ and is generated by differences of these points.

Proof. We use the smooth projective model of C obtained by setting $u_0 = 1$ and eliminating x . Theorem 7.1 is equivalent to the statement on the $\mathbb{Q}(t)$ -points of C : the illegal extension by zero gives rise to the 16 points $(1 : \pm 1 : \pm 1 : \pm 1 : \pm 1)$, and the regular extension by f gives rise to a second orbit of 16 points. By the theorem, these are all the $\mathbb{Q}(t)$ -points on C .

Let E_0, \dots, E_4 be the elliptic curves obtained by eliminating one of the variables u_0, \dots, u_4 . They are given as intersections of two quadrics in \mathbb{P}^3 ; choosing the image of the point $(1 : 1 : 1 : 1 : 1)$ as the origin, we obtain isomorphisms with the curve E considered above and the four further elliptic curves considered in Proposition 6.3 (in a different order). Let $G \subset J(\mathbb{Q}(t))$ denote the subgroup generated by the differences of the $\mathbb{Q}(t)$ -points on C . We find that the image in $E_0 \times \dots \times E_4$ of G is a finite-index subgroup of $G' = (E_0 \times \dots \times E_4)(\mathbb{Q}(t))$ that is isomorphic to \mathbb{Z}^8 ; the index is a power of 2. We know that the kernel of this map (which is the isogeny ϕ) is isomorphic to $(\mathbb{Z}/2\mathbb{Z})^5$ and is contained in the group generated by differences of the $\mathbb{Q}(t)$ -points on C . (This is because the divisors D_{ij} considered in Section 2 can be realized as sums of four $\mathbb{Q}(t)$ -points on C , by pulling back the image of any such

point on Q_{ij} .) It therefore suffices to show that G is 2-saturated. Any element of G that is divisible by 2 maps to an element of G' that is divisible by 2. There are 15 nontrivial cosets of the intersection of the image of G with $2G'$ inside the image of G . We find a representative $P \in G$ of each of these cosets; it is then sufficient to show that no point of the form $P + T$ is divisible by 2, where T is any element of the kernel.

Since we can easily verify that an element of the group of $\mathbb{Q}(t)$ -rational points on the Jacobian of a genus 2 curve is not divisible by 2 by checking that its image under the ‘ $x - T$ map’ is nontrivial (see for example [Sto01]), we consider some of the genus 2 curves D that occur as quotients of C under a subgroup of the automorphism group consisting of sign changes on an even number of variables out of some three-element set. (Such a quotient was also used above in Section 4.) We note that the image of $P + T$ as above must be the sum of the image of P and a $\mathbb{Q}(t)$ -rational 2-torsion point on the Jacobian of D . For each of the 15 choices of P , we find some curve D such that none of these points is divisible by 2 in $\text{Jac}(D)(\mathbb{Q}(t))$. This finishes the proof. \square

REFERENCES

- [BD] Jennifer Balakrishnan and Netan Dogra, *Quadratic Chabauty and rational points I: p -adic heights*, Duke Math. J., available at <https://arxiv.org/abs/1601.00388>. To appear. \uparrow 4.2
- [BD17] ———, *Quadratic Chabauty and rational points II: Generalised height functions on Selmer varieties*, May 1, 2017. Preprint, [arXiv:1705.00401](https://arxiv.org/abs/1705.00401). \uparrow 4.2
- [BCP97] Wieb Bosma, John Cannon, and Catherine Playoust, *The Magma algebra system. I. The user language*, J. Symbolic Comput. **24** (1997), no. 3-4, 235–265, DOI 10.1006/jsco.1996.0125. Computational algebra and number theory (London, 1993). MR1484478 \uparrow 1, 3, 4, 5
- [Bre97] Andrew Bremner, *Some special curves of genus 5*, Acta Arith. **79** (1997), no. 1, 41–51. MR1438115 \uparrow 2
- [Bru03] Nils Bruin, *Chabauty methods using elliptic curves*, J. Reine Angew. Math. **562** (2003), 27–49, DOI 10.1515/crll.2003.076. MR2011330 (2004j:11051) \uparrow 1, 2, 2
- [BS09] Nils Bruin and Michael Stoll, *Two-cover descent on hyperelliptic curves*, Math. Comp. **78** (2009), no. 268, 2347–2370, DOI 10.1090/S0025-5718-09-02255-8. MR2521292 (2010e:11059) \uparrow 2, 4
- [Duj97] Andrej Dujella, *On Diophantine quintuples*, Acta Arith. **81** (1997), no. 1, 69–79, DOI 10.4064/aa-81-1-69-79. MR1454157 \uparrow 3
- [Duj00] ———, *A parametric family of elliptic curves*, Acta Arith. **94** (2000), no. 1, 87–101. MR1762457 \uparrow 1, 6.2
- [Duj04] ———, *There are only finitely many Diophantine quintuples*, J. Reine Angew. Math. **566** (2004), 183–214, DOI 10.1515/crll.2004.003. MR2039327 \uparrow 1
- [Duj] ———, *Diophantine m -tuples web page*. <https://web.math.pmf.unizg.hr/~duje/dtuples.html>. \uparrow 1
- [DGT15] Andrej Dujella, Ivica Gusić, and Petra Tadić, *The rank and generators of Kihara’s elliptic curve with torsion $\mathbf{Z}/4\mathbf{Z}$ over $\mathbf{Q}(t)$* , Proc. Japan Acad. Ser. A Math. Sci. **91** (2015), no. 8, 105–109, DOI 10.3792/pjaa.91.105. MR3403940 \uparrow 6
- [DKMS17] Andrej Dujella, Matija Kazalicki, Miljen Mikić, and Márton Szikszai, *There are infinitely many rational Diophantine sextuples*, Int. Math. Res. Not. IMRN **2** (2017), 490–508, DOI 10.1093/imrn/rnv376. MR3658142 \uparrow 1
- [DPT16] Andrej Dujella, Juan Carlos Peral, and Petra Tadić, *Elliptic curves with torsion group $\mathbf{Z}/6\mathbf{Z}$* , Glas. Mat. Ser. III **51(71)** (2016), no. 2, 321–333, DOI 10.3336/gm.51.2.03. MR3580201 \uparrow 6
- [Fal83] G. Faltings, *Endlichkeitssätze für abelsche Varietäten über Zahlkörpern*, Invent. Math. **73** (1983), no. 3, 349–366 (German). Erratum in: Invent. Math. **75** (1984), 381. MR718935 (85g:11026a) \uparrow 1

- [GJ15] Enrique González-Jiménez, *Covering techniques and rational points on some genus 5 curves*, Fifth Spanish Meeting on Number Theory, Contemp. Math., vol. 649, Amer. Math. Soc., Providence, RI, 2015, pp. 89–105, DOI 10.1090/conm/649/13021. ↑1, 2, 2
- [GT12] Ivica Gusić and Petra Tadić, *A remark on the injectivity of the specialization homomorphism*, Glas. Mat. Ser. III **47(67)** (2012), no. 2, 265–275, DOI 10.3336/gm.47.2.03. MR3006625 ↑1, 5
- [GT15] Ivica Gusić and Petra Tadić, *Injectivity of the specialization homomorphism of elliptic curves*, J. Number Theory **148** (2015), 137–152, DOI 10.1016/j.jnt.2014.09.023. MR3283172 ↑1, 5, 5, 5.3
- [HTZ16] Bo He, Alain Togbé, and Volker Ziegler, *There is no Diophantine quintuple*, October 13, 2016. Preprint, arXiv:1610.04020. ↑1
- [LN59] S. Lang and A. Néron, *Rational points of abelian varieties over function fields*, Amer. J. Math. **81** (1959), 95–118, DOI 10.2307/2372851. MR0102520 ↑5
- [Naj09] Filip Najman, *Integer points on two families of elliptic curves*, Publ. Math. Debrecen **75** (2009), no. 3-4, 401–418. MR2588214 ↑6
- [Shi99] Tetsuji Shioda, *Mordell-Weil lattices for higher genus fibration over a curve*, New trends in algebraic geometry (Warwick, 1996), London Math. Soc. Lecture Note Ser., vol. 264, Cambridge Univ. Press, Cambridge, 1999, pp. 359–373, DOI 10.1017/CBO9780511721540.015. MR1714831 ↑6.2
- [Sil83] Joseph H. Silverman, *Heights and the specialization map for families of abelian varieties*, J. Reine Angew. Math. **342** (1983), 197–211, DOI 10.1515/crll.1983.342.197. MR703488 ↑5
- [Sto01] Michael Stoll, *Implementing 2-descent for Jacobians of hyperelliptic curves*, Acta Arith. **98** (2001), no. 3, 245–277, DOI 10.4064/aa98-3-4. MR1829626 ↑7
- [Sto11] ———, *Rational points on curves*, J. Théor. Nombres Bordeaux **23** (2011), no. 1, 257–277 (English, with English and French summaries). MR2780629 ↑2
- [Sto18] ———, *Chabauty without the Mordell-Weil group*, Algorithmic and Experimental Methods in Algebra, Geometry, and Number Theory (G. Böckle, W. Decker, and G. Malle, eds.), Springer Verlag, 2018, available at <https://arxiv.org/abs/1506.04286>. To appear. ↑2
- [Sto17] ———, *Magma scripts implementing the algorithms in this paper*, 2017. <http://www.mathe2.uni-bayreuth.de/stoll/magma/index.html>. ↑3, 5

MATHEMATISCHES INSTITUT, UNIVERSITÄT BAYREUTH, 95440 BAYREUTH, GERMANY.

E-mail address: Michael.Stoll@uni-bayreuth.de

URL: <http://www.computeralgebra.uni-bayreuth.de>