THE GENERALIZED FERMAT EQUATION $x^2 + y^3 = z^{25}$

NUNO FREITAS AND MICHAEL STOLL

ABSTRACT. We consider the generalized Fermat equation (*) $x^2 + y^3 = z^{25}$. Using the known parameterization of the primitive integral solutions to $x^2 + y^3 = z^5$ (due to Edwards), we reduce the solution of (*) to the solution of five specific equations of the form $H(u, v) = w^5$, where H is homogeneous of degree 10 with coefficients in a sextic number field K, u and v are coprime (rational) integers, and $w \in K$.

1 Introduction

The Generalized Fermat Equation

 $(1.1) x² + y³ = zⁿ, n \ge 6$

is conjectured to have only finitely many non-trivial primitive solutions $(a, b, c) \in \mathbb{Z}^3$ when the "Catalan" solutions $(\pm 3)^2 + (-2)^3 = 1^n$ are counted only once; here *non-trivial* means $abc \neq 0$ and *primitive* means that gcd(a, b, c) = 1. (This is a special case of the more general conjecture that there are only finitely many non-trivial primitive solutions to all equations $x^p + y^q = z^r$ with $1/p + 1/q + 1/r \le 1$.)

Since any integer $n \ge 6$ is divisible by 6, 8, 9, 10, 15, 25 or a prime $p \ge 7$, to study equation (1.1) it suffices to consider these exponents. The cases n = 6, 7, 8, 9, 10 and 15 have been fully resolved and the case n = 11 has been solved assuming GRH. For a more detailed discussion of (1.1), including a list of the known solutions and references to all solved cases, we refer the reader to the introduction of [FNS20].

In this paper we focus on the last remaining non-prime case for n, that is, the equation

(1.2)
$$x^2 + y^3 = z^{25}$$
.

We base our attempt at solving (1.2) on the complete parameterization of the coprime solutions to $x^2 + y^3 = z^5$ obtained by Edwards in [Edw04]. More precisely, using this parameterization, we reduce the resolution of (1.2) to the resolution of several related equations that we proceed to solve using different methods to compute rational points. Unfortunately, we are left with five equations that are out of reach of current methods and computational resources even when assuming GRH. Our main result is the following.

Theorem 1.1. If five specific equations of the form

$$\mathsf{H}(\mathfrak{u},\mathfrak{v})=\mathfrak{w}^5\,,$$

where H is homogeneous of degree 10 with coefficients in a sextic number field K (depending on the equation), u and v are coprime (rational) integers and $w \in K$, have only the expected solutions (see Table 1 below), then the only primitive integer solutions to (1.2)

²⁰¹⁰ Mathematics Subject Classification. Primary 11G30, Secondary 11G35, 14K20.

Key words and phrases. Hyperelliptic curves, descent, Fermat-Catalan, generalized Fermat equation, Selmer set.

are the trivial solutions $(\pm 1, -1, 0)$, $(\pm 1, 0, 1)$, (0, 1, 1), (0, -1, -1) and the non-trivial Catalan solutions $(\pm 3, -2, 1)$.

The assumption above can be verified (or shown to be false) if we are able to determine the sets of L-rational points on five genus 2 curves given by an equation of the form

$$Y^2 = X^5 + A$$

(with $A \in L$), where L is a number field of degree 12 (depending on the equation; it is a quadratic extension of K above).

After recalling Edwards' parameterization in Section 2, which leads to the consideration of 27 equations of the form $z^5 = -h_i(u, v)$, where h_i is a binary form of degree 12, we determine in Section 3 the various possible factorization types of h_i over \mathbb{Q} and over $\mathbb{Q}(\sqrt{5})$. We then eliminate some forms by local considerations; see Section 4. In Section 5 we determine which of the forms in Edwards' list give rise to points on which twist of X(5); these twists correspond to either symplectic or anti-symplectic 5-congruences with an elliptic curve from the list of seven curves obtained in [FNS20].

In the remaining Sections 6, 7, 8 and 9, we deal with the various factorization types in turn. In all cases but the last (when h_i is irreducible over $\mathbb{Q}(\sqrt{5})$), we are able to determine all solutions to (1.2) arising from each form h_i of the corresponding factorization type, using a variety of methods. Some of the last set of forms can also be excluded, and the remaining ones lead to the equations mentioned in Theorem 1.1 above.

In terms of the classification obtained in Section 5, our result is summarized in Table 1. We give the corresponding curve from the list in [FNS20] (where "reducible" refers to the case of reducible 5-torsion) and the symplectic type as "+" or "-". In the equations $H_i(u, v) = w^5$ below, i is the number of an Edwards form, H_i is a binary form of degree 10 with coefficients in a sextic number field K depending on i, u and v are coprime (rational) integers, and $w \in K$.

curve	solutions to (1.2)	condition		
reducible	$(\pm 1, -1, 0)$			
27a1+	$(\pm 1, 0, 1)$			
54a1-	—	$H_{22}(\mathfrak{u},\mathfrak{v})=\mathfrak{w}^5\implies (\mathfrak{u},\mathfrak{v})=(\pm 1,0)$		
96a1+	—	$H_6(\mathfrak{u},\mathfrak{v}) = \mathfrak{w}^5 \implies (\mathfrak{u},\mathfrak{v}) = (\mathfrak{0},\pm 1)$		
288a1+	$\pm(0, 1, 1)$			
864a1+	—	$H_{24}(\mathfrak{u},\mathfrak{v})=\mathfrak{w}^5\implies (\mathfrak{u},\mathfrak{v})=(\pm 1,0)$		
864a1-	—			
864b1+	$(\pm 3, -2, 1)$	$H_5(\mathfrak{u},\mathfrak{v}) = \mathfrak{w}^5 \implies (\mathfrak{u},\mathfrak{v}) = (\mathfrak{0},\pm 1)$		
864b1—	—			
864c1+	—	$H_{16}(u,v) = w^5 \implies (u,v) = (0,\pm 1)$		
864c1-	—			

TABLE 1. Summary of the main result.

The number fields K are given in terms of the minimal polynomial of a generator in Table 2. How the forms H_i can be obtained is explained in Section 9.

i	minimal polynomial
5	$x^6 + 10x^3 + 24x + 5$
6	$x^6 - 2x^5 - 6x - 3$
16	$x^6 + 10x^3 - 15x^2 + 18x - 10$
22	$x^6 + 3x^5 - 10x^3 + 12x - 4$
24	$x^6 - 10x^3 - 6x + 5$

TABLE 2. Coefficient fields of the degree 10 forms.

A Magma script verifying the computational claims in the paper is available on GitHub at [Sto25].

Acknowledgments

We used the Magma Computer Algebra System [BCP97] for the computations.

We thank the Max-Planck-Institut für Mathematik in Bonn for its hospitality on several occasions, which enabled the authors to collaborate on this paper.

2 The Edwards Parameterization

We base our attempt at solving the equation $x^2 + y^3 = z^{25}$ in coprime integers on the complete parameterization of the coprime solutions to $x^2 + y^3 = z^5$ obtained by Edwards in [Edw04], which we now quickly summarize.

In the following, the notation $h = [\alpha_0, \alpha_1, \dots, \alpha_{12}]$ means that h is the binary form

$$h(u,v) = \sum_{i=0}^{12} {12 \choose i} \alpha_i u^i v^{12-i}.$$

We define binary forms h_1, \ldots, h_{27} as given in Table 3.

For $i \in \{1, \ldots, 27\}$, let

$$g_{i} = \frac{1}{132^{2}} \left(\frac{\partial^{2}h_{i}}{\partial u^{2}} \frac{\partial^{2}h_{i}}{\partial v^{2}} - \frac{\partial^{2}h_{i}}{\partial u \partial v} \frac{\partial^{2}h_{i}}{\partial u \partial v} \right) \quad \text{and} \quad f_{i} = \frac{1}{240} \left(\frac{\partial h_{i}}{\partial u} \frac{\partial g_{i}}{\partial v} - \frac{\partial h_{i}}{\partial v} \frac{\partial g_{i}}{\partial u} \right) \,.$$

Note that f_i , g_i and h_i are binary forms with integral coefficients, of degrees 30, 20 and 12, respectively.

Theorem 2.1 (Edwards [Edw04, pages 235–236]). Suppose a, b, c are coprime rational integers satisfying $a^2+b^3+c^5=0$. Then for some $i \in \{1, ..., 27\}$, there is a pair of coprime rational integers u, v and a choice of sign \pm such that

$$a = \pm f_i(u, v)$$
, $b = g_i(u, v)$, $c = h_i(u, v)$.

Suppose that we have a primitive solution (x, y, z) to equation (1.2). Then Theorem 2.1 implies that there are coprime integers u and v satisfying

(2.1)
$$C_i: z^5 = -h_i(u, v)$$

Thus, we have reduced the initial problem to the determination of the triples (u, v, z), with u and v coprime, satisfying one of the 27 possible equations C_i .

 $h_1 = [0, 1, 0, 0, 0, 0, -144/7, 0, 0, 0, 0, -20736, 0]$ $h_2 = [-1, 0, 0, -2, 0, 0, 80/7, 0, 0, 640, 0, 0, -102400]$ $h_3 = [-1, 0, -1, 0, 3, 0, 45/7, 0, 135, 0, -2025, 0, -91125]$ $h_4 = [1, 0, -1, 0, -3, 0, 45/7, 0, -135, 0, -2025, 0, 91125]$ $h_5 = [-1, 1, 1, 1, -1, 5, -25/7, -35, -65, -215, 1025, -7975, -57025]$ $h_6 = [3, 1, -2, 0, -4, -4, 24/7, 16, -80, -48, -928, -2176, 27072]$ $h_7 = [-10, 1, 4, 7, 2, 5, 80/7, -5, -50, -215, -100, -625, -10150]$ $h_8 = [-19, -5, -8, -2, 8, 8, 80/7, 16, 64, 64, -256, -640, -5632]$ $h_{9} = [-7, -22, -13, -6, -3, -6, -207/7, -54, -63, -54, 27, 1242, 4293]$ $h_{10} = [-25, 0, 0, -10, 0, 0, 80/7, 0, 0, 128, 0, 0, -4096]$ $h_{11} = [6, -31, -32, -24, -16, -8, -144/7, -64, -128, -192, -256, 256, 3072]$ $h_{12} = [-64, -32, -32, -32, -16, 8, 248/7, 64, 124, 262, 374, 122, -2353]$ $h_{13} = [-64, -64, -32, -16, -16, -32, -424/7, -76, -68, -28, 134, 859, 2207]$ $h_{14} = [-25, -50, -25, -10, -5, -10, -235/7, -50, -49, -34, 31, 614, 1763]$ $h_{15} = [55, 29, -7, -3, -9, -15, -81/7, 9, -9, -27, -135, -459, 567]$ $h_{16} = [-81, -27, -27, -27, -9, 9, 171/7, 33, 63, 141, 149, -67, -1657]$ $h_{17} = [-125, 0, -25, 0, 15, 0, 45/7, 0, 27, 0, -81, 0, -729]$ $h_{18} = [125, 0, -25, 0, -15, 0, 45/7, 0, -27, 0, -81, 0, 729]$ $h_{19} = [-162, -27, 0, 27, 18, 9, 108/7, 15, 6, -51, -88, -93, -710]$ $h_{20} = [0, 81, 0, 0, 0, 0, -144/7, 0, 0, 0, 0, -256, 0]$ $h_{21} = [-185, -12, 31, 44, 27, 20, 157/7, 12, -17, -76, -105, -148, -701]$ $h_{22} = [100, 125, 50, 15, 0, -15, -270/7, -45, -36, -27, -54, -297, -648]$ $h_{23} = [192, 32, -32, 0, -16, -8, 24/7, 8, -20, -6, -58, -68, 423]$ $h_{24} = [-395, -153, -92, -26, 24, 40, 304/7, 48, 64, 64, 0, -128, -512]$ $h_{25} = [-537, -205, -133, -123, -89, -41, 45/7, 41, 71, 123, 187, 205, -57]$ $h_{26} = [359, 141, -1, -21, -33, -39, -207/7, -9, -9, -27, -81, -189, -81]$ $h_{27} = [295, -17, -55, -25, -25, -5, 31/7, -5, -25, -25, -55, -17, 295]$

TABLE 3. Definition of the forms h_i , $1 \le i \le 27$.

The following observation is important and will be used several times.

Remark 2.2. We note that by construction, all h_i are $GL_2(\mathbb{C})$ -equivalent to the icosahedral Klein form $uv(u^{10} + 11u^5v^5 - v^{10})$. This implies that when K is a number field over which h_i splits off a linear factor, h_i will actually have a factorization of the form

$$h_{i}(u,v) = \ell_{1}(u,v)\ell_{2}(u,v)(A\ell_{1}(u,v)^{10} + B\ell_{1}(u,v)^{5}\ell_{2}(u,v)^{5} + C\ell_{2}(u,v)^{10}),$$

where ℓ_1 and ℓ_2 are linear forms over K and A, B, C \in K (with $(B/11)^2 = AC$). We will make use of this in the following way. Since u and v are coprime, if $-h_i(u, v) = z^5$, we

will have that

$$A\ell_{1}(u,v)^{10} + B\ell_{1}(u,v)^{5}\ell_{2}(u,v)^{5} + C\ell_{2}(u,v)^{10} = \alpha w^{5}$$

for some $\alpha, w \in K$, where α is in a finite set (that has to be determined first). We can write this as either

$$\left(2A\left(\frac{\ell_1(\mathfrak{u},\nu)}{\ell_2(\mathfrak{u},\nu)}\right)^5 + B\right)^2 = 4\alpha A\left(\frac{w}{\ell_2(\mathfrak{u},\nu)^2}\right)^5 + B^2 - 4AC$$

or

$$\left(2C\left(\frac{\ell_2(\mathbf{u},\mathbf{v})}{\ell_1(\mathbf{u},\mathbf{v})}\right)^5 + B\right)^2 = 4\alpha C\left(\frac{w}{\ell_1(\mathbf{u},\mathbf{v})^2}\right)^5 + B^2 - 4AC$$

So a solution gives rise to K-rational points on two genus 2 curves over K,

$$Y^{2} = 4\alpha AX^{5} + (B^{2} - 4AC)$$
 and $Y^{2} = 4\alpha CX^{5} + (B^{2} - 4AC)$

where the points are obtained from

$$X = \frac{w}{\ell_2(u,v)^2}, \quad Y = 2A \left(\frac{\ell_1(u,v)}{\ell_2(u,v)}\right)^5 + B$$

for the first curve, and similarly (with (ℓ_1, A) swapped with (ℓ_2, C)) for the second curve. (One can scale the X and Y coordinates to obtain a curve of the form $Y^2 = X^5 + \gamma$ if desired.)

Note that both curves will be conjugate over the field over which the product $\ell_1 \ell_2$ can be defined when ℓ_1 and ℓ_2 are not yet defined over the base field. So in this case they will provide the same information.

3 Factorization Types

Let $G \in \mathbb{Q}[u, v]$ be a binary form, and let K be a number field. We say G has *factorization type* $[d_1, d_2, \ldots, d_n]$ *over* K if it factors as a product $G = G_1G_2 \cdots G_n$, where the binary forms $G_j \in K[u, v]$ are irreducible over K of degree d_j . Table 4 records the factorization types of h_i over \mathbb{Q} or $\mathbb{Q}(\sqrt{5})$ for $i \in \{1, 2, \ldots, 27\} \setminus \{7, 11, 19\}$. Note that we will show in the next section that the forms associated to $i \in \{7, 11, 19\}$ (which are irreducible over $\mathbb{Q}(\sqrt{5})$) cannot lead to primitive solutions of (1.2).

factorization type of h_i	$i \in I$		
[1, 1, 10] over Q	1, 20, 25		
[4, 8] over Q	3, 4, 12, 17, 18, 27		
[6,6] over $\mathbb{Q}(\sqrt{5})$	2, 10, 26		
[12] over $\mathbb{Q}(\sqrt{5})$	5, 6, 8, 9, 13, 14, 15, 16, 21, 22, 23, 24		

TABLE 4. Factorization types

In the remainder of this work we will study the solutions of (2.1) by applying a strategy that is adapted to each factorization type.

4 Local Primitive Solutions

We can check for which pairs (u, v) of coprime p-adic integers equation C_i above has a solution in \mathbb{Z}_p that comes from a primitive p-adic solution of Equation (1.2). In particular, we find the following.

Lemma 4.1. When $i \in \{7, 11, 19\}$, a solution to equation 1.2 in coprime 2-adic integers cannot give rise to a solution of equation C_i in 2-adic integers.

Proof. This is an easy calculation: for two pairs of coprime residue classes mod 2, the value of $-h_i(u, v)$ has 2-adic valuation 1 and therefore cannot be a fifth power, and for the third coprime residue class, the values of f_i , g_i and h_i are all even, so we do not obtain a primitive solution.

By the same kind of computation, we obtain restrictions on the location of the point $(u : v) \in \mathbb{P}^1(\mathbb{Q}_p)$ for p = 2 and 3. These conditions are recorded in Table 5.

5 The Frey curve and $X_{\rm F}^{\pm}(5)$

As explained in [FNS20, PSS07], to a primitive solution (a, b, c) of the generalized Fermat equation

(5.1) $x^2 + y^3 = z^n$,

where $n \ge 5$ is an integer, we can attach the Frey curve

 $E := E_{(a,b,c)} : y^2 = x^3 + 3bx - 2a.$

We write $\overline{\rho}_{E,p}$ to denote the mod p Galois representation attached to E. The following lemma improves the conclusion of [FNS20, Proposition 2.4] under additional 2-adic and 3-adic hypotheses.

Lemma 5.1. Let (a, b, c) be a primitive solution to (5.1) with exponent $n \ge 7$. Assume at least one of the following conditions:

(i) a *is even or* $b \neq 0, -1, 4 \mod 8$;

(ii) $a \not\equiv \pm 1 \mod 9 \text{ or } b \not\equiv -1 \mod 3$.

Then for all primes $p \ge 5$, the representation $\overline{\rho}_{E,p}$ is absolutely irreducible.

Proof. Observe that if 3 | c, then 3² | 3ⁿ | a² + b³. Therefore, a $\neq \pm 2, \pm 4 \mod 9$ if $b \equiv -1 \mod 3$ as otherwise a² + b³ would be divisible by 3 but not by 9. It follows that (a mod 9, b mod 3) must satisfy one of the congruence conditions in [FNS20, Table 2]. From the proof of this table in *loc. cit.*, we conclude that the Frey curve $E = E_{a,b,c}$ is isomorphic over \mathbb{Q}_3 to a quadratic twist of the curve(s) listed in the fifth column of the same row of the table. Note that except for the curve 96a1, all the curves *W* in the table have conductor N_W satisfying $\nu_3(N_W) = 3$, hence they have semistability defect *e* = 12 and a supercuspidal (hence irreducible) inertial type at 3 (see [DFV24, Table 1]). Since $p \ge 5$ does not divide *e* = 12, we conclude that the inertial subgroup at 3). So $\overline{\rho}_{E,p}$ is irreducible if we avoid 96a1, that is, when a and b satisfy the 3-adic conditions in the statement.

We now apply a similar argument using the Frey curve E/\mathbb{Q}_2 using [FNS20, Table 1]. Indeed, except for the curve 54a1 or the case that E has a quadratic twist with good

	non-excluded residue classes of (u, v)			
i	p = 2 p = 3			
1	(8u, 1)	(81u, 1)		
2	(u, 1)	(3u, 1), (3u + 2, 1), (1, 3v)		
3	(2u, 1), (1, 2v)	(u, 1)		
4	(2u, 1), (1, 2v)	(u, 1)		
5	(2u, 1), (1, 2v)	(3u, 1), (3u + 2, 1), (1, 3v)		
6	(u, 1)	(1, 81v + 51)		
8	(u, 1)	(3u, 1), (3u + 2, 1), (1, 3v)		
9	(2u, 1), (1, 2v)	(u, 1)		
10	(u, 1)	(3u, 1), (3u + 1, 1), (1, 3v)		
12	(1, v)	(3u+2,1), (1,3v), (3u,1)		
13	(1, v)	(3u + 1, 1), (3u, 1), (1, 3v)		
14	(2u, 1), (1, 2v)	(3u + 1, 1), (3u, 1), (1, 3v)		
15	(2u, 1), (1, 2v)	(u, 1)		
16	(2u, 1), (1, 2v)	(3u+2,1), (3u+1,1), (1,3v)		
17	(2u, 1), (1, 2v)	(u, 1)		
18	(2u, 1), (1, 2v)	(u, 1)		
20	(8u, 1)	(1,81v)		
21	(2u, 1), (1, 2v)	(3u+2,1), (3u,1), (1,3v)		
22	(1, 8v + 6)	(u, 1)		
23	(1, v)	(1, 81v + 66)		
24	(u,1)	(3u + 1, 1), (3u, 1), (1, 3v)		
25	(2u, 1), (1, 2v)	(81u + 80, 1)		
26	(2u, 1), (1, 2v)	(u, 1)		
27	(2u, 1), (1, 2v)	(3u+2,1), (3u,1), (1,3v)		

TABLE 5. Residue classes of solutions.

reduction (curve 27a1 or the last row of the table¹) all the curves *W* in the table have semistability defect e = 8 and $v_2(N_W) = 5$. As above, this yields an irreducible inertial type after reduction. We conclude that $\overline{\rho}_{E,p}$ is irreducible if a and b satisfy the 2-adic conditions in the statement.

Our assumptions on (a, b) let us avoid at least one of the two bad cases above, thus $\overline{\rho}_{E,p}$ is irreducible, hence absolutely irreducible since \mathbb{Q} is totally real.

From Table 4 we see that for

 $i \in \mathcal{I} = \{5, 6, 8, 9, 13, 14, 15, 16, 21, 22, 23, 24\}$

¹Note that [FNS20, Table 1] says this case is impossible; however that is proved with a global argument using irreducibility of $\bar{\rho}_{E,p}$ which is what we are trying to prove. For our local argument the last row of the table cannot be excluded.

the corresponding forms are irreducible over \mathbb{Q} and also over $\mathbb{Q}(\sqrt{5})$.

Let now (a, b, c) be a primitive solution to (1.2) arising from a triple $(\pm f_i, g_i, h_i)$ with $i \in \mathcal{I}$, and write E for the associated Frey curve.

For each $i \in \mathcal{I}$ and for all $u, v \in \mathbb{Z}/8\mathbb{Z}$, not both even, we compute the pairs

$$(a,b) = (\pm f_i(u,v) \mod 4, g_i(u,v) \mod 8)$$

and verify in which row (indexed by i_2) of [FNS20, Table 1] it falls. Similarly, for each $i \in \mathcal{I}$ and for all $u, v \in \mathbb{Z}/9\mathbb{Z}$, not both divisible by 3, we compute the pairs

$$(a, b) = (\pm f_i(u, v) \mod 9, g_i(u, v) \mod 3)$$

and verify in which row (indexed by i_3) of [FNS20, Table 2] it falls. Combining this information with [FNS20, Table 3], we can associate to each $i \in \mathcal{I}$ an elliptic curve W_i with Cremona label in {54a1, 96a1, 864a1, 864b1, 864c1}; this correspondence can be found in Table 6. The twists listed in Table 6 come from intersecting the twists in [FNS20, Tables 1 and 2]; note that, for each pair (a, b) as above, the required twist d may differ.

Moreover, the previous calculations also show that the resulting pairs (a, b) always satisfy the hypotheses of Lemma 5.1, hence $\overline{\rho}_{E,5}$ is absolutely irreducible. We now apply [FNS20, Lemma 2.3] to E[5] and conclude that there is a quadratic twist $E^{(d)}$ of E for some $d \in \{\pm 1, \pm 2, \pm 3, \pm 6\}$ such that the the 5-torsion module $E^{(d)}[5]$ is isomorphic to $W_i[5]$ as given by Table 6. Note that [FNS20, Lemma 2.3] as stated applies only to the p-torsion representations $\overline{\rho}_{E_{(a,b,c)},p}$ with $p \ge 7$ a prime, but we can also apply it in our setting. Indeed, we observe that

- (i) the conductor calculations in its proof work for any integer exponent $n \ge 7$ (as used in the proof of Lemma 5.1 above) so, in particular, it holds for our exponent n = 25;
- (ii) if $\overline{\rho}_{E,5}$ is absolutely irreducible, then the level lowering and twisting parts of the argument carry through in the same way.

Since $\overline{\rho}_{E,5}$ has non-abelian image, by [FK22, Theorem 2.1], the symplectic type of the isomorphism $E^{(d)}[5] \simeq W_i[5]$ of $G_{\mathbb{Q}}$ -modules is well defined. That is, either all such isomorphisms are symplectic or all are anti-symplectic (i.e., either all raise the Weil pairing to a power with square exponent or all raise it to a non-square exponent). Thus we obtain a rational point on $X^+_{W_i}(5)$ or $X^-_{W_i}(5)$, respectively, but not on both. To decide which of these is the case for each i, we apply two symplectic criteria.

For i = 22 we have $W_{22} = 54a1$ and both W_{22} and the Frey curve $E = E_{(a,b,c)}^{(d)}$ have multiplicative reduction at 2, so we can apply [FK22, Theorem 1.20]. The minimal discriminants of E and W_{22} are, respectively, $\Delta(E) = 2^{-6}3^3d^6c^{25}$ and $\Delta(W_{22}) = -2^33^9$; thus $v_2(\Delta(E)) \equiv -2v_2(\Delta(W_{22})) \mod 5$, and since -2 is a non-square mod 5, we conclude from [FK22, Theorem 1.20] that the isomorphism $E[5] \simeq W_{22}[5]$ is anti-symplectic. Now observe that, for all $i \neq 22$, the curve W_i has semistability defect e = 8 and conductor N_i with 2-adic valuation $v_2(N_i) = 5$, so we can apply [FK22, Theorem 1.13(A)]. To this end we used a Magma implementation of this theorem. The results can be found in the last column of Table 6.

For completeness we include also in Table 7 the forms that give rise to primitive solutions whose associated Frey curve $E = E_{(a,b,c)}^{(d)}$ is mod 5 congruent to the two CM curves

in the statement of [FNS20, Lemma 2.3], namely 27a1 and 288a1. As above, the irreducibility of $\overline{\rho}_{E,5}$ follows from the inertial types at 2 or 3, and the level lowering and twisting arguments are exactly the same. From [FNS20, Theorem 5.1] we know that we only have to consider the '+' type in all cases; note that although this theorem is stated for $p \ge 11$, the part of its proof required here applies for p = 5 since we know that $\overline{\rho}_{E,5}$ is irreducible.

i	W	d	type
5	864b1	$\pm 2, \pm 6$	+
6	96a1	±3	+
8	864c1	$\pm 2, \pm 6$	_
9	864b1	$\pm 2,\pm 6$	_
13	864b1	$\pm 1, \pm 3$	+
14	864c1	$\pm 1, \pm 3$	+

i	W	d	type
15	864a1	$\pm 2,\pm 6$	—
16	864c1	$\pm 2, \pm 6$	+
21	864a1	$\pm 1, \pm 3$	_
22	54a1	1,-3	—
23	96a1	±6	+
24	864a1	$\pm 2, \pm 6$	+

TABLE 6. Correspondence for $i \in \mathcal{I}$

i	W	d	type
2	27a1	$\pm 2, \pm 6$	+
3	288a1	$\pm 1, \pm 3$	+
4	288a1	$\pm 2, \pm 6$	+
10	27a1	$\pm 2, \pm 6$	+
12	288a1	$\pm 2, \pm 6$	+

i	W	d	type
17	288a1	$\pm 1, \pm 3$	+
18	288a1	$\pm 2, \pm 6$	+
26	27a1	$\pm 2, \pm 6$	+
27	288a1	$\pm 2, \pm 6$	+

TABLE 7. Correspondence to CM curves

6 Factorization type [1, 1, 10] over \mathbb{Q}

Here $i \in \{1, 20, 25\}$ and the polynomial h_i factors over \mathbb{Q} as the product of two linear factors and a factor of degree 10. This is a case where the associated mod 5 representations are reducible.

We apply Remark 2.2 over \mathbb{Q} to the three relevant forms. We have

$$\begin{split} h_1(u,v) &= -12uv(12^4u^{10} + 11 \cdot 12^2u^5v^5 - v^{10}) \\ h_{20}(u,v) &= -12uv(2^8u^{10} + 11 \cdot 12^2u^5v^5 - 3^4v^{10}) \\ h_{25}(u,v) &= -3(u+v)(u-v)\big((u+v)^{10} + 11 \cdot 3^2(u+v)^5(u-v)^5 - 3^4(u-v)^{10}\big) \end{split}$$

Using the coprimality of u and v and the congruence conditions from Table 5, we find that $\alpha = 12, 12, 1$, respectively (where we take $(\ell_1, \ell_2) = (u, v)$ or (u + v, u - v)). The recipe of Remark 2.2 then gives the following pairs of genus 2 curves (after scaling the

coordinates).

$$\begin{array}{lll} \mathfrak{i}=1 & : & Y^2=X^5+2^8\cdot 5^3 & \text{and} & Y^2=X^5+2^6\cdot 3^4\cdot 5^3 \\ \mathfrak{i}=20 & : & Y^2=X^5+2^8\cdot 3^4\cdot 5^3 & \text{and} & Y^2=X^5+2^6\cdot 5^3 \\ \mathfrak{i}=25 & : & Y^2=X^5+2^8\cdot 5^3 & \text{and} & Y^2=X^5+2^8\cdot 3^6\cdot 5^3 \end{array}$$

Using the Magma function MordellWeilGroup, we determine that the group of rational points on the Jacobian variety of the first curve, $Y^2 = X^5 + 2^8 \cdot 5^3$, is infinite cyclic. Then Magma's Chabauty function, applied to a generator of this group, shows that the only rational points on this curve are the point at infinity and a pair of points with x-coordinate -4.

For h_1 , these points correspond to (u, v) = (0, 1) and (0, 1) (one point does not lift), of which the first gives the trivial solution (-1, -1, 0) and the second does not give rise to a primitive solution to the original equation.

For h_{25} , these points correspond to (u, v) = (1, 1) and (1, -1) (again, one point does not lift), both of which do not lead to a primitive solution.

Similarly, we find that the group of rational points on the Jacobian of the curve $Y^2 = X^5 + 2^6 \cdot 5^3$ is trivial, which implies that the point at infinity is the only rational point on that curve. It corresponds to taking (u, v) = (1, 0) in h_{20} , which does not give a primitive solution.

So the only primitive solutions of Equation (1.2) that arise from this case are the trivial solutions $(\pm 1, -1, 0)$.

7 Factorization type [4, 8] over \mathbb{Q}

The mod 5 Galois representation of the associated Frey curves is isomorphic to that of 288a1. Here $i \in \{3, 4, 12, 17, 18, 27\}$ and the polynomial h_i factors over \mathbb{Q} as

$$h_{i}(\mathfrak{u}, \mathfrak{v}) = h_{i,4}(\mathfrak{u}, \mathfrak{v})h_{i,8}(\mathfrak{u}, \mathfrak{v}),$$

where $h_{i,4}$ and $h_{i,8}$ are factors in $\mathbb{Z}[u, v]$ of degree 4 and 8, respectively. With a computer we easily verify the following.

- (1) The prime divisors of the resultant of $h_{i,4}$ and $h_{i,8}$ are 2,3,5. Thus, the prime divisors of $gcd(h_{i,4}(u,v), h_{i,8}(u,v))$ belong to $\{2,3,5\}$ when u and v are coprime integers.
- (2) For p = 2 and p = 3 we see that p does not divide $h_{i,4}(u,v)$ when u and v satisfy the conditions in Table 5.
- (3) Similarly, p = 5 never divides $h_{i,8}(u, v)$ when (u, v) gives rise to a 5-adically primitive solution.

This implies that for coprime $u, v \in \mathbb{Z}$ coming from a primitive solution of (1.2) we always have

$$gcd(h_{i,4}(\mathfrak{u},\mathfrak{v}),h_{i,8}(\mathfrak{u},\mathfrak{v}))=1.$$

It now follows that, if (u, v, z_0) satisfies (2.1) for h_i , then there is an integer z (not divisible by 2 or 3) such that

(7.1)
$$z^5 = h_{i,4}(u,v).$$

We will now show that solutions of (7.1) give rise to rational points on certain hyperelliptic curves. It can be easily checked that, over $\mathbb{Q}(\sqrt{-1})$, we have a factorization

$$h_{i,4}(\mathfrak{u},\mathfrak{v}) = H_i(\mathfrak{u},\mathfrak{v})\bar{H}_i(\mathfrak{u},\mathfrak{v})$$

into conjugate degree two polynomials (see Table 8 for explicit expressions for H_i). Moreover, the resultant of H_i and \bar{H}_i is divisible only by primes above 2 and 3. Since we already know that z above is not divisible by 2 and 3, this implies that $H_i(u,v)$ (and $\bar{H}_i(u,v)$) must be a fifth power in $\mathbb{Z}[\sqrt{-1}]$ (note that all units in $\mathbb{Z}[\sqrt{-1}]$ are fifth powers.) So there are $a, b \in \mathbb{Z}$ such that

(7.2)
$$H_i(u,v) = (a + b\sqrt{-1})^5 = (a^5 - 10a^3b^2 + 5ab^4) + (5a^4b - 10a^2b^3 + b^5)\sqrt{-1}$$
.

Comparing coefficients, this gives

(7.3)
$$a^5 - 10a^3b^2 + 5ab^4 = \operatorname{Re}(H_i(u, v)), \quad 5a^4b - 10a^2b^3 + b^5 = \operatorname{Im}(H_i(u, v)),$$

where Re and Im denote the real and imaginary parts. Now, for each i, we find integers γ_i , α_i , β_i such that

(7.4)
$$\gamma_i \left(\alpha_i \operatorname{Re}(H_i(u, v)) + \beta_i \operatorname{Im}(H_i)(u, v) \right) \operatorname{Im}(H_i(u, v)) = S_i(u, v)^2$$

for some polynomial $S \in \mathbb{Z}[u, v]$. Using (7.3) in (7.4) and writing

(7.5)
$$X = \frac{b}{a}$$
 and $Y = \frac{S_i(u,v)}{a^5}$,

we obtain hyperelliptic curves

$$M_i\colon Y^2=F_i(X)\,,$$

where F_i is of degree 10 and factors as

$$F_i(X) = X(X^4 - 10X^2 + 5)G_i(X)$$

with a polynomial G_i of degree 5. Each solution (u, v) of (2.1) (for one of the indices i considered here) that comes from a primitive solution of (1.2) then gives rise to a rational point on the corresponding curve M_i .

Up to isomorphism, there are only two curves M_i (which are quadratic twists by -1 of each other):

$$\begin{split} F_4(X) &= 3X(X^4 - 10X^2 + 5)(X^5 + 10X^4 - 10X^3 - 20X^2 + 5X + 2) \\ F_{18}(X) &= F_4(-X) \\ F_{12}(X) &= F_{17}(X) = -F_4(X) \\ F_3(X) &= F_{27}(X) = -F_4(-X) \end{split}$$

A summary of the information involved for all i can be found in Table 8.

Note that $(0, 0) \in M_i(\mathbb{Q})$ is a rational point for all i. Applying a partial descent over \mathbb{Q} as in [SS12] to $M_3 \cong M_{12} \cong M_{17} \cong M_{27}$, we obtain a Selmer set with only one element ξ_i , which must correspond to the point (0, 0). Therefore, ξ_i is of the form (?, 5, ?). Thus, given a rational point in $M_i(\mathbb{Q})$, there is a rational point with the same X-coordinate on the genus 1 curve given by

$$C: 5Y^2 = X^4 - 10X^2 + 5$$
.

The Jacobian of C is the elliptic curve

$$E: y^2 = x^3 + x^2 - 83x + 88,$$

i	H _i	γ_i	αί	β_i	$S_i(u,v)$
3	$(3+6\sqrt{-1})u^2+v^2$	3	2	_1	6uv
4	$(-3+6\sqrt{-1})u^2+v^2$	3	2	1	6uv
12	$(2-3\sqrt{-1})u^2 + 2uv + 2v^2$	-3	2	1	3u(u+2v)
17	$3u^2 + (1 - 2\sqrt{-1})v^2$	-3	2	1	6uv
18	$3u^2 + (-1 - 2\sqrt{-1})v^2$	3	-2	1	6uv
27	$(2+\sqrt{-1})u^2 + (2-2\sqrt{-1})uv + (2+\sqrt{-1})v^2$	-3	-2	1	$3(u^2-v^2)$

TABLE 8. Data for the hyperelliptic curves M_i .

which satisfies $E(\mathbb{Q}) \cong \mathbb{Z}/2\mathbb{Z}$. This implies that

$$C(\mathbb{Q}) = \{(0,1), (0,-1)\},\$$

which implies in turn that (0, 0) is the only rational point on M_i .

This argument does not work for $M_4 \cong M_{18}$: M_4 has in addition rational points with $(1,\pm 12)$ (on M_{18} , they are $(-1,\pm 12)$), and the Selmer set correspondingly contains another element (?,-1,?). Unfortunately, the Jacobian elliptic curve of

$$Y^2 = -(X^4 - 10X^2 + 5)$$

has positive rank, so we cannot conclude in the same way. On the other hand, a 2descent on the Jacobian of M_4 shows that the Mordell-Weil rank is at most 1, and the difference P = [(1, 12) - (0, 0)] has infinite order, so the rank is exactly 1 and P generates a subgroup of finite index. We can therefore use Chabauty's method to determine $M_4(\mathbb{Q})$; see below. The result is that

$$M_4(\mathbb{Q}) = \{(0,0), (1,\pm 12)\}$$
 and so $M_{18}(\mathbb{Q}) = \{(0,0), (-1,\pm 12)\}$

Consider a rational point $(b/a, S_i(u, v)/a^5) \in M_i(\mathbb{Q})$. From (7.3), the expression for H_i and the fact that u and v are coprime, we deduce that a and b are coprime (no non-trivial fifth power of an integer can divide both $\text{Re}(H_i(u, v))$ and $\text{Im}(H_i(u, v))$).

We now determine the primitive solutions to equation (1.2) arising from $(0, 0) \in M_i(\mathbb{Q})$. Then b = 0 and $a = \pm 1$, so

$$\operatorname{Re}(\operatorname{H}_{i}(\mathfrak{u},\mathfrak{v})) = \pm 1$$
 and $\operatorname{Im}(\operatorname{H}_{i}(\mathfrak{u},\mathfrak{v})) = 0$.

For $i \in \{12, 17, 17, 27\}$, this pair of equations has no integral solution, and for $i \in \{3, 4\}$, the only solutions are $(u, v) = (0, \pm 1)$ or $(\pm 1, 0)$. These give rise to the trivial solutions (0, 1, 1), (0, -1, -1) and some further non-primitive solutions with a = 0 to equation (1.2).

We now consider the points $(1, \pm 12) \in M_4(\mathbb{Q})$. By the above, we have $a = b = \pm 1$. Using this in (7.3), we obtain the impossible equation $\pm 4 = 6u^2$.

Similarly, the points $(-1,\pm 12) \in M_{18}(\mathbb{Q})$ lead to $a = -b = \pm 1$ and thence to the impossible equation $\pm 4 = -2\nu^2$.

So the only primitive solutions of (1.2) arising from this factorization pattern are $\pm(0, 1, 1)$.

It remains to carry out the Chabauty argument for M_4 .

We use the odd degree model

$$C: y^2 = 30x^9 + 75x^8 - 360x^7 - 300x^6 + 756x^5 + 330x^4 - 360x^3 - 60x^2 + 30x + 30x^4 - 360x^3 - 60x^2 + 30x^4 - 360x^4 - 300x^4 - 360x^4 - 30x^4 -$$

of M_4 , which is obtained by replacing x with 1/x; our goal is then to show that the rational points of C are the point at infinity and the two points $(1, \pm 12)$. Let J denote the Jacobian variety of C.

We work over \mathbb{Q}_{11} since $J(\mathbb{F}_{11}) \cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/70\mathbb{Z} \times \mathbb{Z}/70\mathbb{Z}$ has relatively small exponent. Let $P = [(1, 12) - \infty] \in J(\mathbb{Q})$ be the point of infinite order mentioned above. Then 70P is in the kernel of reduction mod 11. By a standard computation, the details of which we omit here as similar computations can be found in the literature (see also the Magma code at [Sto25]), we determine that the image in $\Omega^1(C/\mathbb{F}_{11})$ of the space of differentials in $\Omega^1(J/\mathbb{Q}_{11})$ that annihilate P is generated by

$$\frac{x-3}{2y} dx, \qquad \frac{x^2-1}{2y} dx \qquad \text{and} \qquad \frac{x^3}{2y} dx$$

The last one of these does not vanish at the point at infinity or at the points $(1, \pm 1)$ in $C(\mathbb{F}_{11})$. From the computation of the 2-Selmer set of C we know that each rational point on C must differ from ∞ or (1, 12) by twice an element of $J(\mathbb{Q})$. Checking this condition mod 11 shows that this implies that every rational point on C reduces mod 11 to either ∞ or a point with x-coordinate 1. The non-vanishing of the reduction of some annihilating differential on these residue classes implies that there can only be one rational point in each of the residue classes, which tells us that there cannot be further rational points.

8 Factorization type [6, 6] over $\mathbb{Q}(\sqrt{5})$

The mod 5 Galois representation of the associated Frey curves is isomorphic to that of 27a1. Here $i \in \{2, 10, 26\}$. The argument in this section is similar to the one used in section 7.

We observe that

$$\begin{aligned} -h_2(-u/2,v) &= -h_{10}(v/2,u) = -h_{26}((u+v)/2,(u-v)/2) \\ &= 25u^{12} + 275u^9v^3 - 165u^6v^6 - 55u^3v^9 + v^{12} \\ &=: h(u,v) \,. \end{aligned}$$

A solution (u, v) in coprime integers to any of the three relevant equations will result in a solution (-2u, v), (2v, u) or (u + v, u - v) in coprime integers of $z^5 = h(u, v)$ (taking into account the restrictions modulo 2 from Table 5).

The polynomial h(u, v) is a quartic in u^3 and v^3 that factors over $\mathbb{Q}(\sqrt{5})$ into two conjugate quadratics in u^3 and v^3 :

$$h(u,v) = \left(v^6 - \frac{55 + 27\sqrt{5}}{2}u^3v^3 - 5u^6\right) \left(v^6 - \frac{55 - 27\sqrt{5}}{2}u^3v^3 - 5u^6\right).$$

The resultant of the two factors is $-3^{18}\sqrt{5}^{12}$, so for coprime integers u and v, the gcd of the two factors on the right hand side is of the form $3^e\sqrt{5}^{e'}$, which implies that each factor is a unit times $3^e\sqrt{5}^{e'}$ times a fifth power. We can assume that $e, e' \in \{0, 1, \ldots, 4\}$. Since the product, which is $3^{2e}5^{e'}$ times the fifth power of a rational integer, must be a

fifth power, it follows that e = e' = 0. So there is some $\alpha \in \mathbb{Z}[\epsilon]$, where $\epsilon = (1 + \sqrt{5})/2$, and some $j \in \{-2, -1, 0, 1, 2\}$ such that

$$\varepsilon^{j}\alpha^{5} = \left(\nu^{6} - \frac{55 + 27\sqrt{5}}{2}u^{3}\nu^{3} - 5u^{6}\right).$$

Writing $\alpha = a + b\sqrt{5}$, we see that

$$\alpha^5 = (a^5 + 50a^3b^2 + 125ab^4) + (5a^4b + 50a^2b^3 + 25b^5)\sqrt{5}.$$

Comparing coefficients, we obtain for j = 0 that

$$a^{5} + 50a^{3}b^{2} + 125ab^{4} = v^{6} - \frac{55}{2}u^{3}v^{3} - 5u^{6}$$
 and $5a^{4}b + 50a^{2}b^{3} + 25b^{5} = \frac{27}{2}u^{3}v^{3}$

Taking appropriate $\mathbb{Q}(\sqrt{-5})$ -linear combinations, we then have

$$(a^{5} + 50a^{3}b^{2} + 125ab^{4}) - \frac{55 \pm 4\sqrt{-5}}{27}(5a^{4}b + 50a^{2}b^{3} + 25b^{5}) = (v^{3} \pm \sqrt{-5}u^{3})^{2}.$$

We take the product of these two (and multiply by 9²) to finally obtain

$$\begin{split} (9(v^6+5u^6))^2 &= 81a^{10} - 1650a^9b + 16725a^8b^2 - 99000a^7b^3 + 395250a^6b^4 - 1039500a^5b^5 \\ &\quad + 1961250a^4b^6 - 2475000a^3b^7 + 2128125a^2b^8 - 1031250ab^9 + 215625b^{10} \\ &=: F(a,b) \,. \end{split}$$

This shows that a solution will lead to a rational point on the hyperelliptic curve of genus 4 given by D_0 : $y^2 = F(x, 1)$, with x = a/b and $y = 9(v^6 + 5u^6)/b^5$.

When $j \neq 0$, we have to expand instead $\varepsilon^{j}(a + b\sqrt{5})^{5} = g_{1}(a, b) + g_{2}(a, b)\sqrt{5}$ with homogeneous polynomials $g_{1}, g_{2} \in \mathbb{Z}[a, b]$ of degree 5 and then perform a similar computation to arrive at another hyperelliptic curve D_{j} of genus 4.

We will see below that $D_2(\mathbb{Q}) = D_1(\mathbb{Q}) = \emptyset$, that $D_0(\mathbb{Q})$ consists of the two points at infinity, and that $D_{-1}(\mathbb{Q})$ and $D_{-2}(\mathbb{Q})$ consist of the two points with x-coordinate 1 (and $y = \pm 192$ and ± 96 , respectively).

Note that the resultant of the two quadratic forms $V^2 - \frac{55\pm27\sqrt{5}}{2}UV - 5U^2$ is $-3^6\sqrt{5}^4$, which implies (since 3^5 is the maximal fifth power of a rational integer that can divide their gcd when evaluated at the coprime integers u^3 and v^3 ; note that if an integer n divides a and b, then n divides both α and its conjugate, so n^5 divides both factors) that either a, $b \in \mathbb{Z}$ and $gcd(a, b) \mid 3$ or $a, b \in \frac{1}{2} + \mathbb{Z}$ and $gcd(2a, 2b) \mid 3$. So we either have b = 0 (from the point at infinity of D_0) and $a \in \{\pm 1, \pm 3\}$ or $a = b \in \{\pm \frac{1}{2}, \pm \frac{3}{2}, \pm 1, \pm 3\}$ (from the points with x = 1 on D_{-1} and D_{-2}). Taking into account that u and v are integers, this gives that

$$v^6 + 5u^6 \in \{1, 3^5, 3^4, 2 \cdot 3^4, 2^5 \cdot 3^4, 2^6 \cdot 3^4\}.$$

The only solutions in coprime integers are $(u, v) = (0, \pm 1)$; they correspond to the trivial solution (1, 0, 1).

So the only primitive solutions to (1.2) arising from this case are the trivial solutions $(\pm 1, 0, 1)$.

We write J_j for the Jacobian of D_j . Using Magma, we can easily check the following facts about the curves D_j .

(1) For j = 1, 2 the routine TwoCoverDescent returns empty fake Selmer sets. Thus $D_1(\mathbb{Q})$ and $D_2(\mathbb{Q})$ are both empty.

- (2) A small search using RationalPoints finds the points $(1 : \pm 9 : 0)$ on D₀, $(1, \pm 192)$ on D₋₁ and $(1, \pm 96)$ on D₋₂.
- (3) The routine TorsionBound tells us that $J_i(\mathbb{Q})$ is torsion free for j = 0, -1, -2.
- (4) The routine RankBound gives a bound of 2 for the rank of the Mordell-Weil groups of J_{-1} and J_{-2} . By taking the difference of the two known points, we get a point in the Jacobian of infinite order. Thus $1 \leq \operatorname{rank}(J_i(\mathbb{Q})) \leq 2$ for j = -1, -2.
- (5) The routine RankBound gives a bound of 4 for the rank of the Mordell-Weil group of J_0 . Again, taking the difference of the two known points we get a point of infinite order. Thus $1 \leq \operatorname{rank}(J_0(\mathbb{Q})) \leq 4$.

In Theorem 9.1 of [Sto17], it is shown that $D_0(\mathbb{Q}) = \{(1 : \pm 9 : 0)\}$ using "Elliptic curve Selmer group Chabauty" (assuming GRH for the necessary class group computations).

It remains to show that D_{-1} and D_{-2} have only the points mentioned earlier. These curves are hyperelliptic of genus 4 and the Mordell-Weil rank of their Jacobians is at most 2. Thus we can hope to find all rational points by applying Chabauty's method. This can be done provided that we can find generators of a subgroup of finite index of $J_i(\mathbb{Q})$.

We first write down the following nicer models for D₁ (which are obtained as

 $SimplifiedModel(ReducedMinimalWeierstrassModel(D_i))$

in Magma).

$$\begin{split} \tilde{D}_{-1}: \quad Y^2 &= 36X^{10} - 120X^9 + 705X^8 - 60X^7 + 3060X^6 + 3846X^5 \\ &\quad + 6390X^4 + 5340X^3 + 3345X^2 + 1230X + 189 \\ \tilde{D}_{-2}: \quad Y^2 &= 9X^{10} - 30X^9 + 645X^8 + 1860X^7 + 6390X^6 + 11274X^5 + 15660X^4 \\ &\quad + 14460X^3 + 8805X^2 + 3120X + 516 \end{split}$$

We write \tilde{J}_j for their Jacobians. A small search for rational points reveals only the points at infinity on both curves. These points were expected to be found since they are the images of the points computed in the previous section under the isomorphism between the models. For both curves we will denote by ∞_+ and ∞_- the points at infinity with positive and negative y-coordinate, respectively. The objective of the remainder of this section is to show that these two points are indeed all the points in $\tilde{D}_j(\mathbb{Q})$.

Lemma 8.1. Let $P \in \tilde{D}_{i}(\mathbb{Q})$. Then the divisor class $[P - \infty_{-}]$ is in $2\tilde{J}_{i}(\mathbb{Q})$.

Proof. Applying the routine TwoCoverDescent to \tilde{D}_j returns a *fake 2-Selmer set* with only one element. By construction we have a factorization of the degree 10 polynomials defining \tilde{D}_j into two conjugate polynomials over $\mathbb{Q}(\sqrt{-5})$. This implies that the fake 2-Selmer set equals the 2-Selmer set. Let C be the unique 2-covering of \tilde{D}_j that has points everywhere locally. Any rational point $P \in \tilde{D}_j(\mathbb{Q})$ lifts to a rational point on some covering having points everywhere locally. Since C is unique, both ∞_- and P lift to a rational point on C. This implies that $[P - \infty_-] \in 2\tilde{J}_j(\mathbb{Q})$.

A computer search reveals the following points $Q_{j,1}, Q_{j,2} \in \tilde{J}_j(\mathbb{Q})$ given in Mumford notation.

$$\begin{aligned} & Q_{-1,1} = (x^4 + 2x^3 + 4x^2 + 3x + 1, \ -60x^3 - 90x^2 - 90x - 30) \\ & Q_{-1,2} = (x^4 - \frac{53}{27}x^3 + \frac{4}{9}x + \frac{1}{9}, \ -\frac{52693}{243}x^3 + \frac{448}{3}x^2 + \frac{8063}{81}x + \frac{1118}{81}) \\ & Q_{-2,1} = (x^4 + 2x^3 + 4x^2 + 3x + 1, \ -30x^3 - 45x^2 - 45x - 15) \\ & Q_{-2,2} = (x^4 + \frac{1}{5}x^3 + \frac{23}{5}x^2 + \frac{21}{5}x + \frac{3}{5}, \ -\frac{683}{25}x^3 + \frac{321}{25}x^2 + \frac{1982}{25}x + \frac{666}{25}) \end{aligned}$$

We note also that $2Q_{j,1} = [\infty_+ - \infty_-]$.

Lemma 8.2. The group $\tilde{J}_{j}(\mathbb{Q})$ is torsion-free and of rank 2. The points $Q_{j,1}$ and $Q_{j,2}$ generate a subgroup $G_{j} \subseteq \tilde{J}_{j}(\mathbb{Q})$ of finite odd index.

Proof. We already know that that $\tilde{J}_{j}(\mathbb{Q})$ is torsion-free and of rank at most 2. We check that in both cases, the image of the subgroup generated by $Q_{j,1}$ and $Q_{j,2}$ under $\tilde{J}_{j}(\mathbb{Q}) \rightarrow \tilde{J}_{j}(\mathbb{F}_{29}) \rightarrow \tilde{J}_{j}(\mathbb{F}_{29})/2\tilde{J}_{j}(\mathbb{F}_{29})$ is isomorphic to $(\mathbb{Z}/2\mathbb{Z})^{2}$. This implies the second claim. \Box

We now want to apply the Chabauty-Coleman method to prove that ∞_{\pm} are the only rational points on \tilde{D}_j . We will use the prime p = 7. Write ρ_7 for the mod 7 reduction maps $\tilde{J}_j(\mathbb{Q}) \to \tilde{J}_j(\mathbb{F}_7)$ and $\tilde{D}_j(\mathbb{Q}) \to \tilde{D}_j(\mathbb{F}_7)$. Denote by ι the Abel-Jacobi embeddings

 $\iota\colon \tilde{D}_j\to \tilde{J}_j \quad \text{given by} \quad \iota(P)=[P-\infty_-]\,.$

Proposition 8.3. Let $G_j = \langle Q_{j,1}, Q_{j,2} \rangle$ as above. Then $\rho_7(G_j) = \rho_7(\tilde{J}_j(\mathbb{Q}))$. Furthermore,

(1) If $P \in \tilde{D}_{-1}(\mathbb{Q})$, then $\rho_7(P) \in \{(1:1:0), (1:6:0), (0:0:1), (1:0:1)\};$ (2) if $P \in \tilde{D}_{-2}(\mathbb{Q})$, then $\rho_7(P) \in \{(1:3:0), (1:4:0)\}.$

Proof. The group $\rho_7(G_j)$ has index 2 in $\tilde{J}_j(\mathbb{F}_7)$. Since G_j has odd index in $\tilde{J}_j(\mathbb{Q})$, we conclude that $\rho_7(G_j) = \rho_7(\tilde{J}_j(\mathbb{Q}))$.

Now for the second statement. Let $P \in \tilde{D}_j(\mathbb{Q})$. From Lemma 8.1 we know that $\iota(P) \in 2\tilde{J}_j(\mathbb{Q})$. Thus, we also have $\rho_7(\iota(P)) \in \rho_7(2\tilde{J}_j(\mathbb{Q}))$. With the computer we easily check that the points $P \in \tilde{D}_j(\mathbb{F}_7)$ such that $[P - \rho_7(\infty_-)] \in \rho_7(2\tilde{J}_j(\mathbb{Q})) = 2\rho_7(G_j)$ are those listed in the statement.

The embedding ι induces an isomorphism ι^* from the space of regular 1-forms $\Omega(\tilde{J}_j/\mathbb{Q}_7)$ to $\Omega(\tilde{D}_j/\mathbb{Q}_7)$. This isomorphism is independent of the base-point ∞_- in the definition of ι . Furthermore, there is a well defined integration pairing

$$\Omega(\tilde{D}_j/\mathbb{Q}_7)\times \tilde{J}_j(\mathbb{Q}_7)\to \mathbb{Q}_7\,,\qquad (\omega,D)\mapsto \int_0^D \iota^{*-1}\omega\,.$$

We need to compute a basis for the differentials that annihilate $\tilde{J}_j(\mathbb{Q}) \subset \tilde{J}_j(\mathbb{Q}_7)$ under this pairing. Since the genus of \tilde{D}_j is 4 and the Mordell-Weil rank of $\tilde{J}_j(\mathbb{Q})$ is 2, we will find two differentials in such a basis. We will now sketch how to obtain this basis of differentials.

We first find two independent points $R_{j,1}$, $R_{j,2}$ in the intersection of $\tilde{J}_j(\mathbb{Q})$ and the kernel of reduction mod 7. We do this because integrals on the kernel of reduction can be computed via power series. We then compute the integrals $\int_0^{R_{j,k}} \iota^{*-1} \omega$ for ω running

through a basis of $\Omega(\tilde{D}_j/\mathbb{Q})$ to sufficient 7-adic precision and finally find the two basis differentials by linear algebra. (We omit the details, as similar computations are described elsewhere in the literature.)

For j = -1, the differentials in the basis reduce mod 7 to

(8.1)
$$\frac{x^3 + 5x^2}{2y} dx \quad \text{and} \quad \frac{x+4}{2y} dx,$$

and for j = -2 the reductions mod 7 are

(8.2)
$$\frac{x^2}{2y} dx$$
 and $\frac{x+4}{2y} dx$

We now determine the rational points on \tilde{D}_{-1} . We see that the differentials mod 7 above do not both vanish at the points in part (1) of Proposition 8.3. This implies that there is at most one point in $\tilde{D}_{-1}(\mathbb{Q})$ reducing to each of these points. In particular, the points ∞_{\pm} are the only rational points in their residue classes. Now let $P \in \tilde{D}_{-1}(\mathbb{F}_7)$ be (0:0:1) or (1:0:1). Suppose there is a rational point W = (x:y:z) reducing to P such that $y \neq 0$. Then (x:-y:z) is another rational point reducing to P. But this is not possible since there can be at most one rational point in these residue classes. Since there is no rational point with vanishing y-coordinate in these residue classes, we conclude that there cannot be any other rational points.

We now determine the rational points on \tilde{D}_{-2} . We consider the points in part (2) of Proposition 8.3. They are the images of the two rational points at infinity. The first of the two differentials mod 7 given above vanishes there to first order. This implies that there are at most two rational points on \tilde{D}_{-2} belonging to each of these two residue classes. To show that there is in fact only one rational point, we take a closer look at the power series expansion of the integral of the second differential at a point at infinity in terms of a uniformizer t there. We obtain

$$I(t) = (13 \cdot 7 + O(7^3))t - (19 \cdot 7 + O(7^3))t^2 - (103 + O(7^3))t^3 + O(t^4).$$

Let $0 \neq a \in \mathbb{Z}_7$. Then (taking into account that the coefficient of t^m in I(t) has valuation at least $-\nu_7(m)$) it follows that

(8.3)
$$\frac{I(7a)}{7^2a} = 13 + O(7^2) - (19 \cdot 7 + O(7^3))a - (103 \cdot 7 + O(7^4))a^2 + O(7^2)$$

which is always $\equiv 6 \mod 7$, so can never vanish. This shows that t = 0 is the only zero of I(t) in $7\mathbb{Z}_7$, and so the point at infinity must be the only rational point in its residue class.

This concludes the proof of

 $D_{-1}(\mathbb{Q}) = \{(1, -192), (1, 192)\}$ and $D_{-2}(\mathbb{Q}) = \{(1, -96), (1, 96)\}$

(for the original models).

9 Forms that are irreducible over $\mathbb{Q}(\sqrt{5})$

The remaining forms we have to deal with are those which are not only irreducible over \mathbb{Q} but also over $\mathbb{Q}(\sqrt{5})$, that is $i \in \mathcal{I} = \{5, 6, 8, 9, 13, 14, 15, 16, 21, 22, 23, 24\}$.

In all cases, there is a sextic number field K_i (that only depends on the mod 5 representation of the associated Frey curves) such that h_i splits as a quadratic form times

a degree 10 form H_i with coefficients in the ring of integers \mathcal{O}_{K_i} of K_i ; we choose the scaling so that H_i is primitive (i.e., its coefficients generate the unit ideal of \mathcal{O}_{K_i}).

The resultant of the two factors is divisible only by primes above 2, 3 and 5 with only one prime above 5 occurring. For $i \notin \{6, 22, 23\}$, we know that c is not divisible by 2 or 3, where $c^5 = -h_i(u, v)$ for coprime integers u and v, so $H_i(u, v)$ cannot be divisible by primes above 2 and 3. For $i \in \{6, 22, 23\}$, we can show (using the congruence conditions from Table 5) that any prime above 2 or 3 must show up with an exponent that is a multiple of 5. The same holds for the possible prime above 5 (which in fact cannot show up). Since all the relevant sextic fields have trivial class group, this implies that a primitive solution to our original equation (1.2) that comes from one of the form considered here must lead to integers u and v such that $H_i(u, v)$ equals a unit times a fifth power in \mathcal{O}_{K_i} . We note that this property is invariant under simultaneous scaling of u and v by a nonzero factor (as that results in multiplying $H_i(u, v)$ by the tenth power of the scaling factor).

We now use a sieving procedure to reduce the number of units (which come from a set of representatives of $\mathcal{O}_{K_i}^{\times}$ modulo fifth powers; the fields K_i have signature (2, 2), so the unit rank is 3, and we have a priori $5^3 = 125$ different units to consider). If p > 5 is a prime, we can run through representatives of all nonzero pairs $(u, v) \in \mathbb{F}_p^2$ modulo scaling and check which units are compatible with the resulting value $H_i(u, v) \in \mathcal{O}_{K_i}/p\mathcal{O}_{K_i} \cong \prod_{\mathfrak{p}|\mathfrak{p}} \mathcal{O}_{K_i}/\mathfrak{p}$. We can also run a similar computation working modulo 25 (which actually leads to quite strong restrictions).

The result of this computation is that *all* units can be excluded when $i \in \{8, 9, 15, 21\}$ (these are the forms that correspond to *antisymplectic* congruences with one of 854a1, 864b1 or 964c1), and in all other cases, only one unit remains. We replace H_i by H_i divided by the corresponding unit, so that the relevant equation is now

(9.1)
$$H_i(u, v) = w^5$$
 with $u, v \in \mathbb{Z}$ and $w \in K_i$.

When we write below "... gives the Frey curve E", this means that E can be obtained as a quadratic twist of the Frey curve associated to the given pair (u, v) and index i. Except for the form with i = 5, the solutions of (9.1) listed below do not give rise to primitive solutions of the original equation (1.2).

- i = 22: (u, v) = (1, 0) gives the Frey curve 54a2 and leads to a solution of (9.1).
- i = 6,23: (u,v) = (0,1) gives the Frey curve 96a1 and leads to a solution of (9.1) (the equations are equivalent).
- i = 24: (u, v) = (1, 0) gives the Frey curve 864a1 and leads to a solution of (9.1).
- i = 5, 13: (u, v) = (0, 1) gives the Frey curve 864b1 and leads to a solution of (9.1) (the equations are equivalent). This gives rise to the primitive pair of solutions $(\pm 3, -2, 1)$ of (1.2).
- i = 14, 16: (u, v) = (1, -1) (for i = 14) and (u, v) = (0, 1) (for i = 16) lead to the Frey curve 864c1 and to a solution of (9.1) (the equations are equivalent).

Proposition 9.1. If the solutions of (9.1) listed above (for $i \in \{5, 6, 13, 14, 16, 22, 23, 24\}$) are the only ones, then the Catalan solutions $(\pm 3, -2, 1)$ are the only primitive solutions to the original equation (1.2) that arise from forms that are irreducible over $\mathbb{Q}(\sqrt{5})$.

Proof. Under the assumption made, all primitive solutions to the original equation arising from one of the remaining eight forms of degree 12 must have (u, v) as in the list

above (up to a common sign change). The only primitive solution that is obtained when evaluating the triples of Edwards forms at these values is the Catalan solution. \Box

Taking into account the fact that three pairs among the eight relevant equations (9.1) are equivalent, and so only five equations need to be considered, this proves the first statement in Theorem 1.1 (given that we have already dealt with all other factorization patterns).

This raises the question how one could try to establish the assumption in the proposition above. Note that equation (9.1) really describes a curve that is given by six equations of the form

 $(\text{degree 10 in } \mathbf{u}, \mathbf{v}) = (\text{quintic in } w_1, \dots, w_6),$

which are obtained by fixing an integral basis of \mathcal{O}_{K_i} , writing w as a linear combination of this basis with coefficients w_1, \ldots, w_6 and then comparing coefficients with respect to this basis on both sides. This is a curve of *very* large genus.

However, one could try to work with the curve over K_i (which by Riemann-Hurwitz has genus 16) and use the extra condition that $u, v \in \mathbb{Z}$ at a suitable stage. But this still seems to be infeasible.

An alternative is to make a further quadratic extension from K_i to L_i such that the quadratic factor in the original factorization of h_i over K_i splits over L_i . Then we can use Remark 2.2 to obtain a genus 2 curve over L_i of the form $Y^2 = X^5 + A$. This proves the second statement in Theorem 1.1. But if we want to use standard methods to try and determine the L_i -rational points on it, we first need to compute its 2-Selmer group, which requires information on the class and unit groups of the field $L_i(\sqrt[5]{A})$ of degree 60 over \mathbb{Q} . Unfortunately, this is quite a bit too large for current technology to give a result in reasonable time, even assuming GRH.

References

- [BCP97] Wieb Bosma, John Cannon, and Catherine Playoust, *The Magma algebra system. I. The user language*, J. Symbolic Comput. **24** (1997), no. 3-4, 235–265, DOI 10.1006/jsco.1996.0125. Computational algebra and number theory (London, 1993). MR1484478 ↑1
- [DFV24] Lassina Dembélé, Nuno Freitaas, and John Voight, On Galois inertial types of elliptic curves over Q_ℓ, April 8, 2024. https://arxiv.org/abs/2203.07787v3, to appear in Math. Comp. ⁵
- [Edw04] Johnny Edwards, *A complete solution to* X² + Y³ + Z⁵ = 0, J. Reine Angew. Math. **571** (2004), 213–236, DOI 10.1515/crll.2004.043. MR2070150 ↑1, 2, 2.1
- [FNS20] Nuno Freitas, Bartosz Naskręcki, and Michael Stoll, *The generalized Fermat equation with exponents* 2, 3, n, Compos. Math. **156** (2020), no. 1, 77–113, DOI 10.1112/s0010437x19007693. MR4036449 ↑1, 1, 5, 5, 5, 1, 5
- [FK22] Nuno Freitas and Alain Kraus, On the symplectic type of isomorphisms of the p-torsion of elliptic curves, Mem. Amer. Math. Soc. 277 (2022), no. 1361, v+105, DOI 10.1090/memo/1361. MR4403927 ↑5
- [PSS07] Bjorn Poonen, Edward F. Schaefer, and Michael Stoll, *Twists of* X(7) *and primitive solutions to* $x^2+y^3 = z^7$, Duke Math. J. **137** (2007), no. 1, 103–158, DOI 10.1215/S0012-7094-07-13714-1. MR2309145 ↑5
- [SS12] Samir Siksek and Michael Stoll, Partial descent on hyperelliptic curves and the generalized Fermat equation x³ + y⁴ + z⁵ = 0, Bull. Lond. Math. Soc. 44 (2012), no. 1, 151–166, DOI 10.1112/blms/bdr086. MR2881333 ↑7
- [Sto25] Michael Stoll, Magma code verifying the computational claims in this paper, 2025. https://
 github.com/MichaelStollBayreuth/Verification. 1, 7
- [Sto17] _____, *Chabauty without the Mordell-Weil group*, Algorithmic and experimental methods in algebra, geometry, and number theory, Springer, Cham, 2017, pp. 623–663. MR3792746 ↑8

INSTITUTO DE CIENCIAS MATEMÁTICAS (ICMAT), C/ NICOLÁS CABRERA 13-15 28049 MADRID, SPAIN *Email address*: nuno.freitas@icmat.es

MATHEMATISCHES INSTITUT, UNIVERSITÄT BAYREUTH, 95440 BAYREUTH, GERMANY *Email address*: Michael.Stoll@uni-bayreuth.de